

# Contractions of 1-loop 5-point tensor Feynman integrals

**Tord Riemann**

DESY, Zeuthen, Germany

in cooperation with **J. Fleischer** (1937-2013), **J. Gluza**, **A. Almasy**

Talk held at Int. Conference

## Matter to the Deepest Recent Developments In Physics of Fundamental Interactions

Ustron, Poland, 1-6 Sep, 2013,

<http://indico.if.us.edu.pl/conferenceDisplay.py?ovw=True&confId=0>






## Definitions

$n$ -point tensor integrals of rank  $R$ :  $(n,R)$ -integrals

$$I_n^{\mu_1 \dots \mu_R} = \int \frac{d^d k}{i\pi^{d/2}} \frac{\prod_{r=1}^R k^{\mu_r}}{\prod_{j=1}^n c_j^{\nu_j}},$$

$d = 4 - 2\epsilon$  and denominators  $c_j$  have *indices*  $\nu_j$  and *chords*  $q_j$

$$c_j = (k - q_j)^2 - m_j^2 + i\epsilon$$



tensor integrals due to, e.g.:

- fermion propagators
- three-gauge boson couplings

## A simple example

### 1-loop self-energy:

$$I_2^\mu = \int \frac{d^d k}{i\pi^{d/2}} \frac{k^\mu}{[k^2 - M_1^2][(k+p)^2 - M_2^2]}$$

**Ansatz :**  $I_2^\mu = p_\mu \cdot B_1(p, M_1, M_2)$

### Solve:

$$\begin{aligned} p_\mu \cdot I_2^\mu &= p^2 \cdot B_1(p, M_1, M_2) \\ &= \int \frac{d^d k}{i\pi^{d/2}} \frac{pk}{[k^2 - M_1^2][(k+p)^2 - M_2^2]} = \int \frac{d^d k}{i\pi^{d/2}} \frac{pk}{D_1 D_2} \\ &= \int \frac{d^d k}{i\pi^{d/2}} \left[ \frac{D_2 - (p^2 - M_2^2 - M_1^2) - D_1}{D_1 D_2} \right], \end{aligned}$$

$$B_1(p, M_1, M_2) = \frac{1}{2p^2} \left[ A_0(M_1) - A_0(M_2) - (p^2 - M_2^2 - M_1^2) B_0(p, M_1, M_2) \right]$$

A **tensor** Feynman integral may be expressed in terms of **scalar** Feynman integrals.

## Passarino-Veltman algorithm

- 1 Contract  $n$ -point and  $R$ -rank Feynman integral with *external momenta*  $p_i^\mu$  and with  $g^{\mu\nu}$ , and cancel propagators
- 2 Invert the resulting system of linear equations
- 3 The result consists of  $(n - 1)$ -point and  $(R - 1)$ -rank functions

Reducing tensor rank introduces inverse Gram determinant:

$$I_5^{\mu_1 \dots \mu_{R-1} \mu_R} = \sum_{i=1}^5 \frac{q_i^{\mu_R}}{\det(G_5)} \left[ A_{0i} I_5^{\mu_1 \dots \mu_{R-1}} - \sum_{s=1}^5 A_{si} I_4^{\mu_1 \dots \mu_{R-1}, s} \right]$$

Gram determinant  $G_n$ :

$$G_n = |2q_i q_j|, \quad i, j = 1, \dots, n-1 \quad (1)$$

and  $A_{0i}$ ,  $A_{si}$  are kinematic coefficients. The  $q_i$  are **internal** momenta.

## Systematic approach to tensor reductions:

1,2,3,4-point functions:

- Passarino, Veltman 1978 [1]

Open-source Source-open programs for 5,6-point reductions:

- LoopTools/FF ( $n \leq 5$ ,  $rank \leq 4$ ), T. Hahn [2, 3] 1998,1990.
- Golem95 T. Binoth et al. [4] 2008
- PJFry ( $n \leq 5$ ,  $rank \leq 5$ ), V. Yundin et al. [5, 6] July 2011

Need in addition a library of scalar functions:

- 't Hooft, Veltman 1979 [7]
- LoopTools/FF T. Hahn [2, 3] 1998,1990
- QCDloop/FF K. Ellis and G. Zanderighi [8, 3] 2007,1990
- OneLooP (complex masses) van Hameren [9] 2010



## History of the Approach - not a complete list of references

- [14] Melrose 1965: Reduction of Feynman diagrams and Cayley determinants
- [15] Davydychev 1991: Integrals in different space-time dimension.
- [16] See also Bern et al. 1993
- [17] Tarasov 1996: Dimensional recurrence relations
- [18] Fleischer, Jegerlehner, Tarasov 2000: 1-loop reductions and signed minors.
  - [4] Binoth, Guillet, Heinrich, Pilon, Schubert, 2005: Algebraic/numerical formalism for one-loop multi-leg amplitudes
- [19] Fleischer and T.Riemann (since 2007) 2011: Complete reduction of 1-loop tensors.
  - See also Diakonidis et al. [20] 2009 and [21] 2009
- [22] Yundin's package PJFry 2011; <https://github.com/Vayu/PJFry>.
  - See also Fleischer, TR, Yundin [5, 6]
- [10] Fleischer and T.Riemann 2011: Contracted tensor Feynman integrals.
- [23] Fleischer and T.Riemann 2012: A solution for tensor reduction of one-loop  $n$ -point functions with  $n \geq 6$



## Tensor integrals expressed in terms of scalar integrals in higher dimensions

$D = d + 2l = 4 - 2\epsilon, 6 - 2\epsilon, \dots$  [Davydychev:1991], also [Fleischer et al.:2000]

$$n_{ij} = \nu_{ij} = 1 + \delta_{ij}, n_{ijk} = \nu_{ij}\nu_{ijk}, \nu_{ijk} = 1 + \delta_{ik} + \delta_{jk}$$

$$I_n^\mu = \int^d k^\mu \prod_{r=1}^n c_r^{-1} = - \sum_{i=1}^n q_i^\mu I_{n,i}^{[d+]}$$

$$I_n^{\mu\nu} = \int^d k^\mu k^\nu \prod_{r=1}^n c_r^{-1} = \sum_{i,j=1}^n q_i^\mu q_j^\nu n_{ij} I_{n,ij}^{[d+]^2} - \frac{1}{2} g^{\mu\nu} I_n^{[d+]}$$

$$I_n^{\mu\nu\lambda} = \int^d k^\mu k^\nu k^\lambda \prod_{r=1}^n c_r^{-1} = - \sum_{i,j,k=1}^n q_i^\mu q_j^\nu q_k^\lambda n_{ijk} I_{n,ijk}^{[d+]^3} + \frac{1}{2} \sum_{i=1}^n g^{\mu\nu} q_i^\lambda I_{n,i}^{[d+]^2}$$

$$\begin{aligned}
 I_n^{\mu\nu\lambda\rho} &= \int \frac{d^d k}{i\pi^{d/2}} \frac{k^\mu k^\nu k^\lambda k^\rho}{\prod_{r=1}^n c_r} = \sum_{i,j,k,l=1}^n q_i^\mu q_j^\nu q_k^\lambda q_l^\rho n_{ijkl} I_{n,ijkl}^{[d+]^4} \\
 &\quad - \frac{1}{2} \sum_{i,j=1}^n g^{\mu\nu} q_i^\lambda q_j^\rho n_{ij} I_{n,ij}^{[d+]^3} + \frac{1}{4} g^{\mu\nu} g^{\lambda\rho} I_n^{[d+]^2}
 \end{aligned} \tag{2}$$

$$\begin{aligned}
I_n^{\mu\nu\lambda\rho\sigma} &= \int \frac{d^d k}{i\pi^{d/2}} \frac{k^\mu k^\nu k^\lambda k^\rho k^\sigma}{\prod_{j=1}^n c_j} \\
&= - \sum_{i,j,k,l,m=1}^n q_i^\mu q_j^\nu q_k^\lambda q_l^\rho q_m^\sigma n_{ijklm} I_{n,ijklm}^{[d+]}{}^5 \\
&\quad + \frac{1}{2} \sum_{i,j,k=1}^n g^{[\mu\nu} q_i^\lambda q_j^\rho q_k^\sigma] n_{ijk} I_{n,ijk}^{[d+]}{}^4 - \frac{1}{4} \sum_{i=1}^n g^{[\mu\nu} g^{\lambda\rho} q_i^\sigma] I_{n,i}^{[d+]}{}^3.
\end{aligned}$$

The integrals are defined in  $[d+]^l = 4 - 2\epsilon + 2l$  dimensions.

$I_{n-1,ab}^{\{\mu_1, \dots\}, s}$ ,  $a, b \neq s$  is obtained from  $I_n^{\{\mu_1, \dots\}}$

by

- shrinking line  $s$
- raising the powers of inverse propagators  $a, b$ .

## Notations: Gram and modified Cayley determinant, signed minors

[Melrose:1965]

Gram determinant  $G_n$ :

$$G_n = |2q_i q_j|, i, j = 1, \dots, n-1 \quad (3)$$

Modified Cayley determinant  $(\ )_N$  of a diagram with  $N$  internal lines and chords  $q_j$ :

$$(\ )_N \equiv \begin{vmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & Y_{11} & Y_{12} & \dots & Y_{1N} \\ 1 & Y_{12} & Y_{22} & \dots & Y_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1N} & Y_{2N} & \dots & Y_{NN} \end{vmatrix} \quad (4)$$

with the matrix elements

$$Y_{ij} = -(q_i - q_j)^2 + m_i^2 + m_j^2, \quad (i, j = 1 \dots N) \quad (5)$$

The propagators are:  $c_i = (k - q_i)^2 - m_i^2$

For the choice  $q_n = 0$ , both determinants are related:

$$(\ )_N = -G_N$$

⇒ The modified Cayley determinant  $(\ )_N$  does not depend on masses.

## Notations: signed minors [Melrose:1965]

signed minors of  $(\ )_N$  are constructed by deleting  $m$  rows and  $m$  columns from  $(\ )_N$ , and multiplying with a sign factor:

$$\begin{aligned} \left( \begin{array}{cccc} j_1 & j_2 & \cdots & j_m \\ k_1 & k_2 & \cdots & k_m \end{array} \right)_N &\equiv \\ &\equiv (-1)^{\sum_i (j_i + k_i)} \operatorname{sgn}_{\{j\}} \operatorname{sgn}_{\{k\}} \left| \begin{array}{c} \text{rows } j_1 \cdots j_m \text{ deleted} \\ \text{columns } k_1 \cdots k_m \text{ deleted} \end{array} \right| \end{aligned} \quad (6)$$

where  $\operatorname{sgn}_{\{j\}}$  and  $\operatorname{sgn}_{\{k\}}$  are the signs of permutations that sort the deleted rows  $j_1 \cdots j_m$  and columns  $k_1 \cdots k_m$  into ascending order.

Example:

$$\left( \begin{array}{c} 0 \\ 0 \end{array} \right)_N \equiv \left| \begin{array}{cccc} Y_{11} & Y_{12} & \cdots & Y_{1N} \\ Y_{12} & Y_{22} & \cdots & Y_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ Y_{1N} & Y_{2N} & \cdots & Y_{NN} \end{array} \right|, \quad (7)$$

## Tarasov's dimensional recurrences for scalars

Following [Tarasov:1996 [17], Fleischer:1999 [18]]  
 apply **recurrence relations**, relating scalar integrals of different dimensions, in order to get rid of the dimensionalities  $[d+]^l = 4 - 2\epsilon + 2l$ :

shift dimension + index:

$$\nu_j(j^+ l_5^{[d+]}) = \frac{1}{(0)_5} \left[ -\binom{j}{0}_5 + \sum_{k=1}^5 \binom{j}{k}_5 \mathbf{k}^- \right] l_5 \quad (8)$$

shift dimension:

$$(d - \sum_{i=1}^5 \nu_i + 1) l_5^{[d+]} = \frac{1}{(0)_5} \left[ \binom{0}{0}_5 - \sum_{k=1}^5 \binom{0}{k}_5 \mathbf{k}^- \right] l_5, \quad (9)$$

shift index:

$$\nu_j j^+ l_5 = \frac{1}{\binom{0}{0}_5} \sum_{k=1}^5 \binom{0j}{0k}_5 \left[ d - \sum_{i=1}^5 \nu_i (\mathbf{k}^- i^+ + 1) \right] l_5 \quad (10)$$

where the operators  $\mathbf{i}^\pm, \mathbf{j}^\pm, \mathbf{k}^\pm$  act by shifting the indices  $\nu_i, \nu_j, \nu_k$  by  $\pm 1$ .

## Recursions for tensors – Alternative to dimensional recurrences of scalars

Express any  $(5, R)$  pentagon by a  $(5, R - 1)$  pentagon plus  $(4, R - 1)$  boxes

[Diakonidis, Fleischer, T. Riemann, Tausk: Phys. Lett. **B683** (2010) [21]]

5-point tensor recursion:

$$I_5^{\mu_1 \dots \mu_{R-1} \mu} = I_5^{\mu_1 \dots \mu_{R-1}} Q_0^\mu - \sum_{s=1}^5 I_4^{\mu_1 \dots \mu_{R-1}, s} Q_s^\mu,$$

For  $n = 6, 7, 8, \dots$  things are close but differ a bit.

auxiliary vectors with inverse Gram determinants

$$Q_s^\mu = \sum_{i=1}^5 q_i^\mu \frac{\binom{s}{i}_5}{\binom{s}{s}_5}, \quad s = 0, \dots, 5$$

For e.g.  $R = 3$ , again  $[1/\binom{s}{i}_5]^3$  will occur.

## Contractions

One may combine now *Tarasov's dimensional recurrence relations* and the *tensor rank recursions* in order to derive especially useful representations.

After that, we will perform contractions with external momenta.

So, the following equations are symbolic:

$$q_{i_1 \mu_1} \cdots q_{i_R \mu_R} I_5^{\mu_1 \cdots \mu_R} = \int \frac{d^d k}{i\pi^{d/2}} \frac{\prod_{r=1}^R (q_{i_r} \cdot k)}{\prod_{j=1}^5 c_j},$$

$$g_{\mu_1, \mu_2} q_{i_1 \mu_3} \cdots q_{i_R \mu_R} I_5^{\mu_1 \cdots \mu_R} \neq \int \frac{k^2 d^d k}{i\pi^{d/2}} \frac{\prod_{r=3}^R (q_{i_r} \cdot k)}{\prod_{j=1}^5 c_j}$$

One may arrange a one-loop calculation such that all the one-loop integrals appear **only** in such contractions.

**Important:**

The contraction with  $g_{\mu_1, \mu_2}$  is shown here in a symbolic form; in practice we work strictly 4-dimensional with  $g_{\mu_1, \mu_2}$ .







As an example, we reproduce the 4-point part of  $I_{4,ijkl}^{[d+]}$ :

$$\begin{aligned}
 n_{ijkl} I_{4,ijkl}^{[d+]} &= \frac{\binom{0}{i} \binom{0}{j} \binom{0}{k} \binom{0}{l}}{\binom{0}{0} \binom{0}{0} \binom{0}{0} \binom{0}{0}} d(d+1)(d+2)(d+3) I_4^{[d+]} \\
 &+ \frac{\binom{0j}{0j} \binom{0k}{0k} \binom{0l}{0l} + \binom{0i}{0k} \binom{0l}{0j} \binom{0l}{0i} + \binom{0j}{0k} \binom{0l}{0i} \binom{0l}{0j} + \binom{0i}{0l} \binom{0l}{0j} \binom{0k}{0k} + \binom{0j}{0l} \binom{0l}{0i} \binom{0k}{0k} + \binom{0k}{0l} \binom{0l}{0i} \binom{0j}{0j}}{\binom{0}{0}^3} \\
 &\times d(d+1) I_4^{[d+]} \\
 &+ \frac{\binom{0i}{0l} \binom{0j}{0k} + \binom{0j}{0l} \binom{0i}{0k} + \binom{0k}{0l} \binom{0i}{0j}}{\binom{0}{0}^2} I_4^{[d+]} + \dots
 \end{aligned} \tag{12}$$

In (12), one has to understand the 4-point integrals to carry the corresponding index  $s$  and the signed minors are  $\binom{0}{k} \rightarrow \binom{0s}{ks}_5$  etc.

✓ no scalar 5-point integrals in higher dimensions

✓ no inverse Gram det.  $\binom{0}{5}$

✓ **4-point integrals without indices**

† scalar 4-point integrals in higher dimensions:  $I_4^{[d+],s}$  etc.

† inverse Gram det.  $\binom{0}{5} \equiv \binom{0}{4}$

## Contractions for the 5-point functions with rank $R = 1$

A chord is the momentum shift of an internal line due to external momenta,  $D_i = (k - q_i)^2 - m_i^2 + i\epsilon$ , and  $q_i = (p_1 + p_2 + \dots + p_i)$ , with  $q_n = 0$ .

The tensor 5-point integral of rank  $R = 1$  is ([19], eq. (4.6)):

$$\begin{aligned} I_5^\mu &= - \sum_{i=1}^5 q_i^\mu I_{5,i}^{[d^+]} \\ &= - \sum_{i=1}^4 q_i^\mu \sum_{s=1}^5 \frac{\binom{0i}{0s}_5}{\binom{0}{0}_5} I_4^s \end{aligned} \quad (13)$$

This yields, when contracted with a chord,

$$q_{a\mu} I_5^\mu = - \frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \left[ \sum_{i=1}^4 (q_a \cdot q_i) \binom{0i}{0s}_5 \right] I_4^s. \quad (14)$$

In fact, the **sum over  $i$  may be performed explicitly**:

$$\Sigma_a^{1,s} \equiv \sum_{i=1}^4 (q_a \cdot q_i) \begin{pmatrix} 0s \\ 0i \end{pmatrix}_5 = +\frac{1}{2} \left\{ \begin{pmatrix} s \\ 0 \end{pmatrix}_5 (Y_{a5} - Y_{55}) + \begin{pmatrix} 0 \\ 0 \end{pmatrix}_5 (\delta_{as} - \delta_{5s}) \right\},$$

We get immediately

$$q_{a\mu} l_5^\mu = -\frac{1}{\begin{pmatrix} 0 \\ 0 \end{pmatrix}_5} \sum_{s=1}^5 \Sigma_a^{1,s} l_4^s. \quad (15)$$

## Contractions for the 5-point functions with rank $R = 2$

$$I_5^{\mu\nu} = \sum_{i,j=1}^4 q_i^\mu q_j^\nu E_{ij} + g^{\mu\nu} E_{00}, \quad (16)$$

has the following tensor coefficients free of  $1/()_5$ :

$$E_{00} = - \sum_{s=1}^5 \frac{1}{2} \frac{1}{\binom{0}{0}_5} \binom{s}{0}_5 I_4^{[d+],s}, \quad (17)$$

$$E_{ij} = \sum_{s=1}^5 \frac{1}{\binom{0}{0}_5} \left[ \binom{0i}{sj}_5 I_4^{[d+],s} + \binom{0s}{0j}_5 I_{4,i}^{[d+],s} \right]. \quad (18)$$

Equation (16) yields for the contractions with chords:

$$q_{a\mu} q_{b\nu} l_5^{\mu\nu} = \sum_{i,j=1}^4 (q_a \cdot q_i)(q_b \cdot q_j) E_{ij} + (q_a \cdot q_b) E_{00}. \quad (19)$$

and finally (19) simply reads

$$\begin{aligned} q_{a\mu} q_{b\nu} l_5^{\mu\nu} &= \frac{1}{4} \sum_{s=1}^5 \left\{ \frac{\binom{s}{0}_5}{\binom{0s}{0s}_5} (\delta_{ab} \delta_{as} + \delta_{5s}) + \frac{\binom{s}{0}_5}{\binom{0s}{0s}_5} [(\delta_{as} - \delta_{5s})(Y_{b5} - Y_{55}) \right. \\ &\quad \left. + (\delta_{bs} - \delta_{5s})(Y_{a5} - Y_{55}) + \frac{\binom{s}{0}_5}{\binom{0}{0}_5} (Y_{a5} - Y_{55})(Y_{b5} - Y_{55})] \right\} l_4^{[d+],s} \\ &\quad + \frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \frac{\sum_b^{1,s}}{\binom{0s}{0s}_5} \sum_{t=1}^5 \sum_a^{2,st} l_3^{st}, \end{aligned}$$

with

$$\begin{aligned}\Sigma_a^{2,st} &\equiv \sum_{i=1}^4 (q_a \cdot q_i) \begin{pmatrix} 0st \\ 0si \end{pmatrix}_5 \\ &= \frac{1}{2} (1 - \delta_{st}) \left\{ \begin{pmatrix} ts \\ 0s \end{pmatrix}_5 (Y_{a5} - Y_{55}) + \begin{pmatrix} 0s \\ 0s \end{pmatrix}_5 (\delta_{at} - \delta_{5t}) - \begin{pmatrix} 0s \\ 0t \end{pmatrix}_5 (\delta_{as} - \delta_{5s}) \right\}\end{aligned}$$

This has been extended also to higher ranks.

We need at most double sums, e.g.:

$$\begin{aligned}\Sigma_{ab}^{2,s} &\equiv \sum_{i,j=1}^4 (q_a \cdot q_i)(q_b \cdot q_j) \begin{pmatrix} si \\ sj \end{pmatrix}_5 \\ &= \frac{1}{2} (q_a \cdot q_b) \begin{pmatrix} s \\ s \end{pmatrix}_5 - \frac{1}{4} ()_5 (\delta_{ab}\delta_{as} + \delta_{5s}), \quad (20)\end{aligned}$$

The **sums over signed minors, weighted with scalar products of chords** are given in J. Fleischer, T.R., PLB 2011 [10].



## Contractions for the 5-point functions with rank $R = 4$

$$I_5^{\mu\nu\lambda\rho} = I_5^{\mu\nu\lambda} \cdot Q_0^\rho - \sum_{s=1}^5 I_4^{\mu\nu\lambda,s} \cdot Q_s^\rho. \quad (21)$$

Contracted with chords (differences of external momenta):

$$q_a^\mu q_b^\nu q_c^\lambda q_d^\rho I_5^{\mu\nu\lambda\rho} = q_a^\mu q_b^\nu q_c^\lambda I_5^{\mu\nu\lambda} (q_d^\rho Q_0^\rho) - C_{5,abcd} \quad (22)$$

Here:

$$Q_s^\rho = \sum_{i=1}^5 q_i^\rho \frac{\binom{s}{i}_5}{\binom{0}{5}}, \quad s = 0 \dots 5 \quad (23)$$

The first term  $q_a^\mu q_b^\nu q_c^\lambda I_5^{\mu\nu\lambda}$  is known, the second term has to be determined:

$$C_{5,abcd} = - \sum_{s=1}^5 q_{a\mu} q_{b\nu} q_{c\lambda} I_4^{\mu\nu\lambda,s} \frac{1}{2} (\delta_{ds} - \delta_{5s}), \quad (24)$$

becomes:

$$\begin{aligned}
C_{5,abcd} = & \frac{1}{16} \left\{ G^5 + \delta_{ab}\delta_{ac}\delta_{ad}G^d - I_1^{5abc} - I_1^{5abd} - I_1^{5acd} - I_1^{5bcd} + I_1^{abcd} - J_3^{a5} - J_3^{b5} - J_3^{c5} - J_3^{d5} \right. \\
& + R^{5ab} + R^{5ac} + R^{5bc} + R^{5da} + R^{5db} + R^{5dc} + \delta_{bc}\delta_{bd} \left( J_3^{ad} - J_3^{5d} \right) + \delta_{ac}\delta_{ad} \left( J_3^{bd} - J_3^{5d} \right) \\
& + \delta_{ab}\delta_{ad} \left( J_3^{cd} - J_3^{5d} \right) + \delta_{ab}\delta_{ac} \left( J_3^{dc} - J_3^{5c} \right) + \delta_{ab}\delta_{cd}\tilde{J}_3^{db} + \delta_{ad}\delta_{bc}\tilde{J}_3^{dc} + \delta_{ac}\delta_{bd}\tilde{J}_3^{dc} \\
& + \delta_{ab} \left( \tilde{J}_3^{5b} - R^{b5c} - R^{bd5} + R^{bdc} \right) + \delta_{ac} \left( \tilde{J}_3^{5c} - R^{c5b} - R^{cd5} + R^{cdb} \right) \\
& + \delta_{ad} \left( \tilde{J}_3^{d5} - R^{d5b} - R^{d5c} + R^{dbc} \right) + \delta_{bc} \left( \tilde{J}_3^{5c} - R^{c5a} - R^{cd5} + R^{cda} \right) \\
& + \delta_{bd} \left( \tilde{J}_3^{d5} - R^{d5a} - R^{d5c} + R^{dac} \right) + \delta_{cd} \left( \tilde{J}_3^{d5} - R^{d5a} - R^{d5b} + R^{dab} \right) \\
& + \delta_{ab}\delta_{ad}J_4^d Y_c + \delta_{ac}\delta_{ad}J_4^d Y_b + \delta_{bc}\delta_{bd}J_4^d Y_a + \delta_{ab} \left( R^{bd} - R^{b5} \right) Y_c + \delta_{ac} \left( R^{cd} - R^{c5} \right) Y_b \\
& + \delta_{ad} \left( R^{dc} - R^{d5} \right) Y_b + \delta_{ad} \left( R^{db} - R^{d5} \right) Y_c + \delta_{bc} \left( R^{cd} - R^{c5} \right) Y_a + \delta_{bd} \left( R^{dc} - R^{d5} \right) Y_a \\
& + \delta_{bd} \left( R^{da} - R^{d5} \right) Y_c + \delta_{cd} \left( R^{da} - R^{d5} \right) Y_b + \delta_{cd} \left( R^{db} - R^{d5} \right) Y_a + \left( I_4^d - I_4^5 \right) Y_a Y_b Y_c \\
& + \left( I_3^{cd} - I_3^{5c} - I_3^{5d} + R^5 + \delta_{cd}R^d \right) Y_a Y_b + \left( I_3^{bd} - I_3^{5b} - I_3^{5d} + R^5 + \delta_{bd}R^d \right) Y_a Y_c \\
& + \left( I_3^{ad} - I_3^{5a} - I_3^{5d} + R^5 + \delta_{ad}R^d \right) Y_b Y_c + \left( I_2^{bcd} - I_2^{5bc} - I_2^{5bd} - I_2^{5cd} - J_4^5 + R^{5b} + R^{5c} + R^{5d} \right) Y_a \\
& + \left( I_2^{acd} - I_2^{5ac} - I_2^{5ad} - I_2^{5cd} - J_4^5 + R^{5a} + R^{5c} + R^{5d} \right) Y_b \\
& + \left. \left( I_2^{abd} - I_2^{5ab} - I_2^{5ad} - I_2^{5bd} - J_4^5 + R^{5a} + R^{5b} + R^{5d} \right) Y_c \right\} \tag{25}
\end{aligned}$$

where we have introduced:

$$J_3^{st} \equiv \frac{1}{\binom{st}{st}_5} \left\{ -\binom{S}{S}_5 I_3^{[d+],st} + \binom{ts}{0S}_5 R^{ts} - \sum_{u=1}^5 \binom{ts}{us}_5 R^{tsu} \right\}, \quad (26)$$

$$\tilde{J}_3^{st} \equiv \frac{1}{\binom{st}{st}_5} \left\{ \binom{S}{t}_5 I_3^{[d+],st} + \binom{St}{0t}_5 R^{ts} - \sum_{u=1}^5 \binom{St}{ut}_5 R^{tsu} \right\}, \quad (27)$$

$$G^s \equiv \frac{1}{\binom{s}{s}_5} \left\{ -2\binom{\quad}{s}_5 R^{[d+],s} + \binom{S}{0}_5 J_4^s - \sum_{t=1}^5 \binom{S}{t}_5 J_3^{ts} \right\}. \quad (28)$$

$J_4^s$  and  $R^{[d+],s}$  are given in Eqs. (2.24) and (2.44) of [19] respectively.

We assume throughout that  $q_5 = 0$ , where  $q_1, \dots, q_5$  are chords – differences of external momenta.

Further abbreviations (see (2.24, 2.49, 2.9, 2.17, 2.34, 2.41) of [19]) :

$$J_4^s \equiv \frac{1}{\binom{s}{5}} \left\{ -\binom{d+1}{5} I_4^{[d+1],s} + \binom{s}{0}_5 R^s - \sum_{t=1}^5 \binom{s}{t}_5 R^{st} \right\} \quad (29)$$

$$= \frac{-1}{\binom{0s}{5}} \left\{ \binom{1}{5} (d-2)(d-1) I_4^{[d+1]^2,s} - \binom{0}{5} I_4^{[d+1],s} + \sum_{t=1}^5 \binom{t}{5} (d-2) I_3^{[d+1],st} + \sum_{t=1}^5 \binom{0s}{0t}_5 R^{st} \right\}$$

$$R^{[d+1],s} = \frac{1}{\binom{s}{5}} \left[ \binom{s}{0}_5 I_4^{[d+1],s} - \sum_{t=1}^5 \binom{s}{t}_5 I_3^{[d+1],st} \right] = \frac{1}{\binom{0s}{5}} \left[ \binom{s}{0}_5 (d-1) I_4^{[d+1]^2,s} - \sum_{t=1}^5 \binom{0s}{0t}_5 I_3^{[d+1],st} \right] \quad (30)$$

$$R^s \equiv \frac{1}{\binom{s}{5}} \left[ \binom{s}{0}_5 I_4^s - \sum_{t=1}^5 \binom{s}{t}_5 I_3^t \right] = \frac{1}{\binom{0s}{5}} \left[ \binom{s}{0}_5 I_4^{[d+1],s} - \sum_{t=1}^5 \binom{0s}{0t}_5 I_3^{st} \right]. \quad (31)$$

$$R^{st} \equiv \frac{1}{\binom{st}{5}} \left[ \binom{st}{0t}_5 I_3^{st} - \sum_{u=1}^5 \binom{st}{ut}_5 I_2^{stu} \right] = \frac{1}{\binom{0st}{5}} \left[ \binom{st}{0t}_5 (d-2) I_3^{[d+1],st} - \sum_{u=1}^5 \binom{0st}{0ut}_5 I_2^{stu} \right]. \quad (32)$$

$$R^{tsu} = \frac{1}{\binom{tsu}{5}} \left[ \binom{tsu}{0su}_5 I_2^{tsu} - \sum_{v=1}^5 \binom{tsu}{vsu}_5 I_1^{stuv} \right]$$

$$= \frac{1}{\binom{0tsu}{5}} \left[ \binom{tsu}{0su}_5 (d-1) I_2^{[d+1],tsu} - \sum_{v=1}^5 \binom{0tsu}{0vsu}_5 I_1^{stuv} \right]. \quad (33)$$

$$Y_a = Y_{a5} - Y_{55}, \quad Y_{ab} = -(q_a - q_b)^2 + m_a^2 + m_b^2 \quad (34)$$

## Contractions for the 5-point functions with rank $R = 5$ – NEW, to be tested yet – I

For the rank  $R = 5$  case  $I_5^{\mu\nu\lambda\rho\sigma}$ , we have three types of contractions:

- contraction with momenta,  $q_{a\mu} q_{b\nu} q_{c\lambda} q_{d\rho} q_{e\sigma}$
- contractions with momenta and **one metric tensor**,  $q_{a\mu} q_{b\nu} q_{c\lambda} g_{\rho\sigma}$
- contractions with momenta and **two metric tensors**,  $q_{a\mu} g_{\nu\lambda} g_{\rho\sigma}$

The contraction with momenta yields the following final expression:

$$\begin{aligned}
 CE5(a, b, c, d, e) &\equiv q_{a\mu} q_{b\nu} q_{c\lambda} q_{d\rho} q_{e\sigma} I_5^{\mu\nu\lambda\rho\sigma} \\
 &= -\frac{1}{2} CE4(a, b, c, d) Y_e + C_{5,abcde}
 \end{aligned} \tag{35}$$

where

$$C_{5,abcde} = - \sum_{s=1}^5 q_{a\mu} q_{b\nu} q_{c\lambda} q_{d\rho} I_4^{\mu\nu\lambda\rho,s} \frac{1}{2} (\delta_{es} - \delta_{5s}),$$

is explicitly evaluated to be:

$$\begin{aligned}
C_{5,abcde} = & \frac{1}{32} \left\{ \delta_{ab}\delta_{ac}\delta_{ad}\delta_{ae}F^e - F^1 + J_2^{5ea} + J_2^{5eb} + J_2^{5ec} + J_2^{ab5} + J_2^{ac5} + J_2^{ad5} + J_2^{bc5} + J_2^{bd5} \right. \\
& + J_2^{cd5} + J_2^{de5} - G^{a5} - G^{b5} - G^{c5} - G^{d5} - G^{e5} - R^{5abc} - R^{5abd} - R^{5abe} - R^{5acd} - R^{5ace} - R^{5ade} \\
& - R^{5bcd} - R^{5bce} - R^{5bde} - R^{5cde} - \delta_{ae}\delta_{bc}\delta_{bd}G_1^{de} - \delta_{ac}\delta_{ad}\delta_{be}G_1^{de} - \delta_{ab}\delta_{ad}\delta_{ce}G_1^{de} \\
& - \delta_{ab}\delta_{cd}\delta_{ce}G_1^{eb} - \delta_{ad}\delta_{ae}\delta_{bc}G_1^{ec} - \delta_{ac}\delta_{bd}\delta_{be}G_1^{ec} + \delta_{bc}\delta_{bd}\delta_{be} \left( G^{5e} - G^{ae} - G^e Y_a \right) \\
& + \delta_{ac}\delta_{ad}\delta_{ae} \left( G^{5e} - G^{be} - G^e Y_b \right) + \delta_{ab}\delta_{ad}\delta_{ae} \left( G^{5e} - G^{ce} - G^e Y_c \right) + \delta_{ab}\delta_{ac}\delta_{ae} \left( G^{5e} - G^{de} - G^e Y_d \right) \\
& + \delta_{ab}\delta_{ac}\delta_{ad} \left( G^{5d} - G^{ed} \right) - \delta_{ac}\delta_{ae}\delta_{bd}\tilde{G}^{de} - \delta_{ad}\delta_{bc}\delta_{be}\tilde{G}^{de} - \delta_{ab}\delta_{ae}\delta_{cd}\tilde{G}^{de} - \delta_{ab}\delta_{ac}\delta_{de}\tilde{G}^{ec} \\
& + \delta_{bc}\delta_{bd} \left[ J_2^{5ad} + J_2^{5ed} - J_2^{aed} - G_1^{d5} + \left( J_3^{5d} - J_3^{ed} \right) Y_a \right] \\
& + \delta_{ac}\delta_{ad} \left[ J_2^{5bd} + J_2^{5ed} - J_2^{bed} - G_1^{d5} + \left( J_3^{5d} - J_3^{ed} \right) Y_b \right] \\
& + \delta_{ab}\delta_{ad} \left[ J_2^{5cd} + J_2^{5ed} - J_2^{ced} - G_1^{d5} + \left( J_3^{5d} - J_3^{ed} \right) Y_c \right] \\
& + \delta_{cd}\delta_{ce} \left[ J_2^{5ae} + J_2^{5be} - J_2^{abe} - G_1^{e5} + \left( J_3^{5e} - J_3^{be} \right) Y_a + \left( J_3^{5e} - J_3^{ae} \right) Y_b - J_4^e Y_a Y_b \right] \\
& + \delta_{bd}\delta_{be} \left[ J_2^{5ae} + J_2^{5ce} - J_2^{ace} - G_1^{e5} + \left( J_3^{5e} - J_3^{ce} \right) Y_a + \left( J_3^{5e} - J_3^{ae} \right) Y_c - J_4^e Y_a Y_c \right] \\
& + \delta_{ad}\delta_{ae} \left[ J_2^{5be} + J_2^{5ce} - J_2^{bce} - G_1^{e5} + \left( J_3^{5e} - J_3^{ce} \right) Y_b + \left( J_3^{5e} - J_3^{be} \right) Y_c - J_4^e Y_b Y_c \right] \\
& + \delta_{bc}\delta_{de} \left( \tilde{J}_2^{5ce} - \tilde{J}_2^{ace} - \tilde{J}_3^{ce} Y_a \right) + \delta_{be}\delta_{cd} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{ade} - \tilde{J}_3^{de} Y_a \right) + \delta_{bd}\delta_{ce} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{ade} - \tilde{J}_3^{de} Y_a \right) \\
& + \delta_{ac}\delta_{de} \left( \tilde{J}_2^{5ce} - \tilde{J}_2^{bce} - \tilde{J}_3^{ce} Y_b \right) + \delta_{ae}\delta_{cd} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{bde} - \tilde{J}_3^{de} Y_b \right) + \delta_{ad}\delta_{ce} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{bde} - \tilde{J}_3^{de} Y_b \right) \\
& + \delta_{ab}\delta_{de} \left( \tilde{J}_2^{5be} - \tilde{J}_2^{cbe} - \tilde{J}_3^{be} Y_c \right) + \delta_{ae}\delta_{bd} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{cde} - \tilde{J}_3^{de} Y_c \right) + \delta_{ad}\delta_{be} \left( \tilde{J}_2^{5de} - \tilde{J}_2^{cde} - \tilde{J}_3^{de} Y_c \right) \\
& + \delta_{ab}\delta_{ce} \left( \tilde{J}_2^{5be} - \tilde{J}_2^{dbe} - \tilde{J}_3^{be} Y_d \right) + \delta_{ae}\delta_{bc} \left( \tilde{J}_2^{5ce} - \tilde{J}_2^{dce} - \tilde{J}_3^{ce} Y_d \right) + \delta_{ac}\delta_{be} \left( \tilde{J}_2^{5ce} - \tilde{J}_2^{dce} - \tilde{J}_3^{ce} Y_d \right) \dots
\end{aligned}$$

$$\begin{aligned}
& \dots + \delta_{ab}\delta_{cd} (\tilde{J}_2^{5bd} - \tilde{J}_2^{ebd}) + \delta_{ad}\delta_{bc} (\tilde{J}_2^{5cd} - \tilde{J}_2^{ecd}) + \delta_{ac}\delta_{bd} (\tilde{J}_2^{5cd} - \tilde{J}_2^{ecd}) \\
& + \delta_{ab}\delta_{ac} \left[ J_2^{5dc} + J_2^{5ec} - J_2^{dec} - \tilde{G}^{5c} + (J_3^{5c} - J_3^{ec}) Y_d \right] \\
& + \delta_{bc}\delta_{be} \left[ J_2^{5ae} + J_2^{5de} - J_2^{ade} - \tilde{G}^{5e} + (J_3^{5e} - J_3^{de}) Y_a + (J_3^{5e} - J_3^{ae}) Y_d - J_4^e Y_a Y_d \right] \\
& + \delta_{ac}\delta_{ae} \left[ J_2^{5be} + J_2^{5de} - J_2^{bde} - \tilde{G}^{5e} + (J_3^{5e} - J_3^{de}) Y_b + (J_3^{5e} - J_3^{be}) Y_d - J_4^e Y_b Y_d \right] \\
& + \delta_{ab}\delta_{ae} \left[ J_2^{5ce} + J_2^{5de} - J_2^{cde} - \tilde{G}^{5e} + (J_3^{5e} - J_3^{de}) Y_c + (J_3^{5e} - J_3^{ce}) Y_d - J_4^e Y_c Y_d \right] \\
& + \delta_{ab} \left[ G_1^{5b} - \tilde{J}_2^{5cb} - \tilde{J}_2^{d5b} - \tilde{J}_2^{e5b} + R^{b5cd} + R^{b5ce} + R^{b5de} - R^{bcde} + (-\tilde{J}_3^{5b} + R^{b5d} + R^{b5e} - R^{bde}) Y_c \right. \\
& \left. + (-\tilde{J}_3^{5b} + R^{b5c} + R^{b5e} - R^{bce}) Y_d + (R^{b5} - R^{be}) Y_c Y_d \right] + \delta_{bc} \left[ G_1^{5c} - \tilde{J}_2^{a5c} - \tilde{J}_2^{d5c} - \tilde{J}_2^{e5c} + R^{c5ad} + R^{c5de} \right. \\
& \left. + R^{c5ae} - R^{cade} - R^{bcde} + (-\tilde{J}_3^{5c} + R^{c5d} + R^{c5e} - R^{cde}) Y_a + (-\tilde{J}_3^{5c} + R^{c5a} + R^{c5e} - R^{cae}) Y_d \right. \\
& \left. + (R^{c5} - R^{ce}) Y_a Y_d \right] + \delta_{ac} \left[ G_1^{5c} - \tilde{J}_2^{b5c} - \tilde{J}_2^{d5c} - \tilde{J}_2^{e5c} + R^{c5bd} + R^{c5be} + R^{c5de} - R^{cbde} \right. \\
& \left. + (-\tilde{J}_3^{5c} + R^{c5d} + R^{c5e} - R^{cde}) Y_b + (-\tilde{J}_3^{5c} + R^{c5b} + R^{c5e} - R^{cbe}) Y_d + (R^{c5} - R^{ce}) Y_b Y_d \right] \\
& + \delta_{ce} \left[ G_1^{5e} - \tilde{J}_2^{a5e} - \tilde{J}_2^{b5e} - \tilde{J}_2^{d5e} + R^{e5ab} + R^{e5ad} + R^{e5bd} - R^{eabd} + (-\tilde{J}_3^{5e} + R^{e5b} + R^{e5d} - R^{ebd}) Y_a \right. \\
& \left. + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5d} - R^{ead}) Y_b + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5b} - R^{eab}) Y_d + (R^{e5} - R^{ed}) Y_a Y_b \right. \\
& \left. + (R^{e5} - R^{eb}) Y_a Y_d + (R^{e5} - R^{ea}) Y_b Y_d - R^e Y_a Y_b Y_d \right] + \delta_{be} \left[ G_1^{5e} - \tilde{J}_2^{a5e} - \tilde{J}_2^{c5e} - \tilde{J}_2^{d5e} + R^{e5ac} + R^{e5ad} \right. \\
& \left. + R^{e5cd} - R^{eacd} + (-\tilde{J}_3^{5e} + R^{e5c} + R^{e5d} - R^{ecd}) Y_a + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5d} - R^{ead}) Y_c \right. \\
& \left. + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5c} - R^{eac}) Y_d + (R^{e5} - R^{ed}) Y_a Y_c + (R^{e5} - R^{ec}) Y_a Y_d + (R^{e5} - R^{ea}) Y_c Y_d - R^e Y_a Y_c Y_d \right] \\
& + \delta_{ae} \left[ G_1^{5e} - \tilde{J}_2^{b5e} - \tilde{J}_2^{c5e} - \tilde{J}_2^{d5e} + R^{e5bc} + R^{e5bd} + R^{e5cd} - R^{ebcd} + (-\tilde{J}_3^{5e} + R^{e5c} + R^{e5d} - R^{ecd}) Y_b \dots \right]
\end{aligned}$$

$$\begin{aligned}
& \dots + (-\tilde{J}_3^{5e} + R^{e5b} + R^{e5d} - R^{ebd}) Y_c + (-\tilde{J}_3^{5e} + R^{e5b} + R^{e5c} - R^{ebc}) Y_d + (R^{e5} - R^{ed}) Y_b Y_c \\
& + (R^{e5} - R^{ec}) Y_b Y_d + (R^{e5} - R^{eb}) Y_c Y_d - R^e Y_b Y_c Y_d + \delta_{cd} [-\tilde{J}_2^{a5d} - \tilde{J}_2^{b5d} - \tilde{J}_2^{e5d} + R^{d5ab} + R^{d5ae} + R^{d5be} \\
& - R^{dabe} + \tilde{G}^{d5} + (-\tilde{J}_3^{d5} + R^{d5b} + R^{d5e} - R^{dbe}) Y_a + (-\tilde{J}_3^{d5} + R^{d5a} + R^{d5e} - R^{dae}) Y_b + (R^{d5} - R^{de}) Y_a Y_b] \\
& + \delta_{bd} [-\tilde{J}_2^{a5d} - \tilde{J}_2^{c5d} - \tilde{J}_2^{e5d} + R^{d5ac} + R^{d5ae} + R^{d5ce} - R^{dace} + \tilde{G}^{d5} + (-\tilde{J}_3^{d5} + R^{d5c} + R^{d5e} - R^{dce}) Y_a \\
& + (-\tilde{J}_3^{d5} + R^{d5a} + R^{d5e} - R^{dae}) Y_c + (R^{d5} - R^{de}) Y_a Y_c] + \delta_{ad} [-\tilde{J}_2^{b5d} - \tilde{J}_2^{c5d} - \tilde{J}_2^{e5d} + R^{d5bc} + R^{d5be} \\
& + R^{d5ce} - R^{dbce} + \tilde{G}^{d5} + (-\tilde{J}_3^{d5} + R^{d5c} + R^{d5e} - R^{dce}) Y_b + (-\tilde{J}_3^{d5} + R^{d5b} + R^{d5e} - R^{dbe}) Y_c \\
& + (R^{d5} - R^{de}) Y_b Y_c] + \delta_{de} [-\tilde{J}_2^{a5e} - \tilde{J}_2^{b5e} - \tilde{J}_2^{c5e} + R^{e5ab} + R^{e5ac} + R^{e5bc} - R^{eabc} + \tilde{G}^{e5} \\
& + (-\tilde{J}_3^{5e} + R^{e5b} + R^{e5c} - R^{ebc}) Y_a + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5c} - R^{eac}) Y_b + (-\tilde{J}_3^{5e} + R^{e5a} + R^{e5b} - R^{eab}) Y_c \\
& + (R^{e5} - R^{ec}) Y_a Y_b + (R^{e5} - R^{eb}) Y_a Y_c + (R^{e5} - R^{ea}) Y_b Y_c - R^e Y_a Y_b Y_c] + (I_1^{5bcd} + I_1^{5bce} + I_1^{5bde} + I_1^{5cde} \\
& - I_1^{bcde} + J_3^{b5} + J_3^{c5} + J_3^{d5} + J_3^{e5} - G^5 - R^{5bc} - R^{5bd} - R^{5be} - R^{5cd} - R^{5ce} - R^{5de}) Y_a + (I_1^{5acd} + I_1^{5ace} \\
& + I_1^{5ade} + I_1^{5cde} - I_1^{acde} + J_3^{a5} + J_3^{c5} + J_3^{d5} + J_3^{e5} - G^5 - R^{5ac} - R^{5ad} - R^{5ae} - R^{5cd} - R^{5ce} - R^{5de}) Y_b \\
& + (I_1^{5abd} + I_1^{5abe} + I_1^{5ade} + I_1^{5bde} - I_1^{abde} + J_3^{a5} + J_3^{b5} + J_3^{d5} + J_3^{e5} - G^5 - R^{5ab} - R^{5ad} - R^{5ae} - R^{5bd} \\
& - R^{5be} - R^{5de}) Y_c + (I_1^{5abc} + I_1^{5abe} + I_1^{5ace} + I_1^{5bce} - I_1^{abce} + J_3^{a5} + J_3^{b5} + J_3^{c5} + J_3^{e5} - G^5 - R^{5ab} \\
& - R^{5ac} - R^{5ae} - R^{5bc} - R^{5be} - R^{5ce}) Y_d + (I_2^{5cd} + I_2^{5ce} + I_2^{5de} - I_2^{cde} + J_4^5 - R^{5c} - R^{5d} - R^{5e}) Y_a Y_b \\
& + (I_2^{5bd} + I_2^{5be} + I_2^{5de} - I_2^{bde} + J_4^5 - R^{5b} - R^{5d} - R^{5e}) Y_a Y_c + (I_2^{5bc} + I_2^{5be} + I_2^{5ce} - I_2^{bce} + J_4^5 - R^{5b} \\
& - R^{5c} - R^{5e}) Y_a Y_d + (I_2^{5ad} + I_2^{5ae} + I_2^{5de} - I_2^{ade} + J_4^5 - R^{5a} - R^{5d} - R^{5e}) Y_b Y_c + (I_2^{5ac} + I_2^{5ae} + I_2^{5ce} \dots
\end{aligned}$$



$$\begin{aligned}
& \dots - I_2^{ace} + J_4^5 - R^{5a} - R^{5c} - R^{5e}) Y_b Y_d + (I_2^{5ab} + I_2^{5ae} + I_2^{5be} - I_2^{abe} + J_4^5 - R^{5a} - R^{5b} - R^{5e}) Y_c Y_d \\
& + (I_3^{5d} + I_3^{5e} - I_3^{de} - R^5) Y_a Y_b Y_c + (I_3^{5c} + I_3^{5e} - I_3^{ce} - R^5) Y_a Y_b Y_d + (I_3^{5b} + I_3^{5e} - I_3^{be} - R^5) Y_a Y_c Y_d \\
& + (I_3^{5a} + I_3^{5e} - I_3^{ae} - R^5) Y_b Y_c Y_d + (I_4^5 - I_4^e) Y_a Y_b Y_c Y_d \}. \tag{36}
\end{aligned}$$

with some additional definitions not shown here

## Some numerics for 5-point function with rank $R = 4$

We use OneLOop for scalar functions and compare with LoopTools/FF

the kinematics:

```

pls = 1.1163688400000000E-002  p2s = 2.6109999999999998E-007  p3s = 0.0000000000000000
p4s = 2.6109999999999998E-007  p5s = 1.1163688400000000E-002
s12 = -0.70858278190000001      s23 = -1.5343299000000002E-003  s34 = -0.12851860429999998
s45 = -0.61023937949999996      s15 = 0.92668942420000000
m1s = 1.1163688361676107E-002  m2s = 0.00000000000000000      m3s = 2.6112003932088364E-007
m4s = 2.6112003932088364E-007  m5s = 0.00000000000000000

```

OneLOop-3.3.1

is used for the evaluation of 1-loop scalar 1-, 2-, 3- and 4-point functions

[van Hameren arXiv:1007.4716] and [van Hameren, Papadopoulos, Pittau arXiv:0903.4665]

```

the R=4 contractions, a,b,c,d=3,3,3,3 ( -48094.1074 54542318 , -47802.08746 5035322 )
LoopTools ( -48094.1074 65 , -47802.08746 05 )

```

```

the R=4 contractions, a,b,c,d=3,3,3,4 ( -18463.1204 24842149 , -23446.4704 12257226 )
LoopTools ( -18463.1204 31 , -23446.4704 09 )

```

```

the R=4 contractions, a,b,c,d=3,3,3,5 ( 0.0000000000000000 , 0.0000000000000000 )
LoopTools ( 0.0000000000000000 , 0.0000000000000000 )

```

compared to: LoopTools/FF [Hahn arXiv:hep-ph/9807565] and [van Oldenborgh CPC66(1991)]

## Summary

- Explicit **Analytical, recursive treatment** of **heptagon, hexagon and pentagon tensor integrals** of rank  $R$  in terms of pentagons and boxes of rank  $R - 1$
- Systematic derivation of expressions which are explicitly **free of inverse Gram determinants**  $(\ )_5$  until pentagons of rank  $R = 5$
- Numerical **tensor reduction package PJFry** for C, C++, Mathematica, Fortran
- Numerical **packages for contracted tensor integrals OLEC and CONTRACTIONS** for C++ and Fortran under development

## References I

- [1] G. Passarino, M. Veltman, One Loop Corrections for  $e^+e^-$  Annihilation Into  $\mu^+\mu^-$  in the Weinberg Model, Nucl. Phys. B160 (1979) 151.  
doi:[10.1016/0550-3213\(79\)90234-7](https://doi.org/10.1016/0550-3213(79)90234-7).
- [2] T. Hahn, M. Perez-Victoria, Automatized one loop calculations in four-dimensions and D-dimensions, Comput. Phys. Commun. 118 (1999) 153–165.  
arXiv:[hep-ph/9807565](https://arxiv.org/abs/hep-ph/9807565), doi:[10.1016/S0010-4655\(98\)00173-8](https://doi.org/10.1016/S0010-4655(98)00173-8).
- [3] G. J. van Oldenborgh, FF: A Package to evaluate one loop Feynman diagrams, Comput. Phys. Commun. 66 (1991) 1–15, doi:[10.1016/0010-4655\(91\)90002-3](https://doi.org/10.1016/0010-4655(91)90002-3), scanned version at [http://ccdb3fs.kek.jp/cgi-bin/img\\_index?9004168](http://ccdb3fs.kek.jp/cgi-bin/img_index?9004168).  
doi:[10.1016/0010-4655\(91\)90002-3](https://doi.org/10.1016/0010-4655(91)90002-3).
- [4] T. Binoth, J. P. Guillet, G. Heinrich, E. Pilon, T. Reiter, Golem95: a numerical program to calculate one-loop tensor integrals with up to six external legs, Comput. Phys. Commun. 180 (2009) 2317–2330.  
arXiv:[0810.0992](https://arxiv.org/abs/0810.0992), doi:[10.1016/j.cpc.2009.06.024](https://doi.org/10.1016/j.cpc.2009.06.024).
- [5] V. Yundin, Massive loop corrections for collider physics, PhD thesis, Humboldt-Universität zu Berlin, 2012, see <http://edoc.hu-berlin.de/docviews/abstract.php?id=39163> and urn:nbn:de:kobv:11-100199626.
- [6] J. Fleischer, T. Riemann, V. Yundin, PJFry: A C++ package for tensor reduction of one-loop Feynman integrals. In: [22], preprint DESY 11-25 (2011), <http://www-library.desy.de/cgi-bin/showprep.pl?desy11-252>.
- [7] G. 't Hooft, M. Veltman, Scalar One Loop Integrals, Nucl. Phys. B153 (1979) 365–401.  
doi:[10.1016/0550-3213\(79\)90605-9](https://doi.org/10.1016/0550-3213(79)90605-9).
- [8] R. K. Ellis, G. Zanderighi, Scalar one-loop integrals for QCD, JHEP 02 (2008) 002.  
arXiv:[arXiv:0712.1851](https://arxiv.org/abs/0712.1851), doi:[10.1088/1126-6708/2008/02/002](https://doi.org/10.1088/1126-6708/2008/02/002).

## References II

- [9] A. van Hameren, OneLOop: For the evaluation of one-loop scalar functions, *Comput. Phys. Commun.* 182 (2011) 2427–2438.  
[arXiv:arXiv:1007.4716](https://arxiv.org/abs/1007.4716), doi:10.1016/j.cpc.2011.06.011.
- [10] J. Fleischer, T. Riemann, Calculating contracted tensor Feynman integrals, *Phys. Lett. B* 701 (2011) 646–653.  
[arXiv:arXiv:1104.4067](https://arxiv.org/abs/1104.4067), doi:10.1016/j.physletb.2011.06.033.
- [11] J. Fleischer, J. Gluza, M. Gluza, and T. Riemann, OLEC project; C++ and Mathematica packages OLEC v.0.9 available at Silesian Univ., Katowice, webpage <http://prac.us.edu.pl/gluza/olec/>.
- [12] A. Almasy, Mathematica and Fortran packages for contracted 1-loop tensor reduction, to be published.
- [13] J. Fleischer, T. Riemann, **A solution for tensor reduction of one-loop N-point functions with  $N \geq 6$** , *Physics Letters B* 707 (2012) 375–380.  
[arXiv:arXiv:1111.5821](https://arxiv.org/abs/1111.5821), doi:10.1016/j.physletb.2011.12.060.  
 URL <http://www.sciencedirect.com/science/article/pii/S0370269311015280>
- [14] D. B. Melrose, Reduction of Feynman diagrams, *Nuovo Cim.* 40 (1965) 181–213.  
 doi:10.1007/BF028329.
- [15] A. I. Davydychev, A simple formula for reducing Feynman diagrams to scalar integrals, *Phys. Lett. B* 263 (1991) 107–111, doi:10.1016/0370-2693(91)91715-8.  
 doi:10.1016/0370-2693(91)91715-8.
- [16] Z. Bern, L. J. Dixon, D. A. Kosower, Dimensionally Regulated One-Loop Integrals, *Phys. Lett. B* 302 (1993) 299–308 [Erratum–*ibid.* B 318 (1993) 649].  
[arXiv:hep-ph/9212308](https://arxiv.org/abs/hep-ph/9212308), doi:10.1016/0370-2693(93)90400-C.

## References III

- [17] O. Tarasov, Connection between Feynman integrals having different values of the space-time dimension, Phys.Rev. D54 (1996) 6479–6490.  
[arXiv:hep-th/9606018](#), doi:10.1103/PhysRevD.54.6479.
- [18] J. Fleischer, F. Jegerlehner, O. Tarasov, Algebraic reduction of one-loop Feynman graph amplitudes, Nucl. Phys. B566 (2000) 423–440.  
[arXiv:hep-ph/9907327](#), doi:10.1016/S0550-3213(99)00678-1.
- [19] J. Fleischer, T. Riemann, Complete algebraic reduction of one-loop tensor Feynman integrals, Phys. Rev. D83 (2011) 073004.  
[arXiv:arXiv:1009.4436](#), doi:10.1103/PhysRevD.83.073004.
- [20] T. Diakonidis, J. Fleischer, J. Gluza, K. Kajda, T. Riemann, J. Tausk, A complete reduction of one-loop tensor 5- and 6-point integrals, Phys. Rev. D80 (2009) 036003.  
[arXiv:arXiv:0812.2134](#), doi:10.1103/PhysRevD.80.036003.
- [21] T. Diakonidis, J. Fleischer, T. Riemann, J. B. Tausk, A recursive reduction of tensor Feynman integrals, Phys. Lett. B683 (2010) 69–74.  
[arXiv:arXiv:0907.2115](#), doi:10.1016/j.physletb.2009.11.049.
- [22] J. Alcaraz Maestre, et al., The SM and NLO Multileg and SM MC Working Groups: Summary Report, report of the SM and NLO Multileg and SM MC Working Groups for the Workshop 'Physics at TeV Colliders', Les Houches, France, 31 May – 8 June, 2011. See [\[arXiv:1203.1488\]](#) and [\[arXiv:1203.6803\]](#) (2012).  
[arXiv:1203.6803](#).
- [23] J. Fleischer, T. Riemann, A solution for tensor reduction of one-loop N-point functions with  $N \geq 6$ , Phys. Lett. B707 (2012) 375–380.  
[arXiv:1111.5821](#), doi:10.1016/j.physletb.2011.12.060.