

”Collider Physics” 2nd Symposium of the Division for Physics of Fundamental Interactions of the Polish Physical Society

Friday, 13 May 2016 - Sunday, 15 May 2016

Katowice

Book of Abstracts

Contents

Dark matter near a resonance	1
Axino dark matter with low reheating temperature	1
4-jet production in high-energy factorization	1
Advances in amplitudes with off-shell partons	2
Clocks and dynamics in Quantum Universe	2
Probing TeV scale physics with neV neutrons	2
Relaxation rates and phase transitions	3
Maximum tension principle and entropic force	3
The evolution of cosmological domain walls from Higgs effective potential	3
Dark matter - how well can one determine its properties?	4
First detection of a gravitational wave signal	4
Recent ATLAS and CMS results on the Standard Model and the Higgs boson	4
Beyond the Standard Model searches in the Higgs sector from ATLAS and CMS	5
Study of matter at extreme conditions with ALICE	5
Recent ATLAS and CMS results on the Standard Model and the Higgs boson	5
Beyond the Standard Model searches in the Higgs sector from ATLAS and CMS	5
Study of matter at extreme conditions with ALICE	5
First detection of a gravitational wave signal	5
Clocks and dynamics in Quantum Universe	6
Dark matter - how well can one determine its properties?	6
Dark matter near a resonance	6
The averaging problem in cosmology	6
Maximum tension principle and entropic force	6

Uncovering BFKL dynamics in production of Mueller-Navelet jets at the LHC	6
Soft gluon resummation in associated Higgs boson production	6
Forward Physics with the ATLAS Detector. Results, Prospects and Polish Contribution .	7
From exclusive production, through diffraction, to jets correlations - forward physics re- sults from CMS	7
Status of exotic four-quark mesons	7
Relaxation rates and phase transitions	7
Effective particles in quantum field theory	7
Lattice QCD: a numerical tool for precision physics in the Standard Model and beyond .	7
Conformal Standard Model	7
The averaging problem in cosmology	8
In search of higher twist effects in proton structure	8
Hints for the BSM Physics at the LHC	8
BSM physics at e+e- colliders	8
Phenomenology of Gauge Mediation models at the LHC and future colliders	8
Enabling Electroweak Baryogenesis through Dark Matter	9
Successes and challenges in perturbative QCD	9
Matching fixed order QCD with parton shower for Drell-Yan and Higgs production . . .	9
Advances in amplitudes with off-shell partons	9
Radiative corrections to hadron production in e+e- annihilation	9
4-jet production in high-energy factorization	9
4-jet production in high-energy factorization	10
Probing TeV scale physics with neV neutrons	10
Recent Belle results	10
Recent results from LHCb	10
Recent results from LHCb	10
Belle II perspective	10
LHCb perspectives on flavour physics	10
Neutrino properties determined in oscillation experiments	11
Astrophysical searches with neutrino detectors	11

Neutrino portal to new physics	11
--	----

0

Dark matter near a resonance

Authors: Bohdan Grządkowski¹; Mateusz Duch²

¹ *University of Warsaw*

² *Faculty of Physics, University of Warsaw*

Corresponding Author: mateusz.duch@fuw.edu.pl

We study dark matter annihilation via a mediator with a mass in the vicinity of the resonance. In this case, Breit-Wigner enhancement leads to the averaged cross-section that strongly depends on the temperature. As a result, the standard "freeze-out" mechanism is altered and an effective annihilation can last long after decoupling from the equilibrium. We show, how a dark matter with the large self-interaction cross-section can be obtained in this framework and comment on the tension with indirect detection experiments, particularly Fermi-LAT observations of dwarf spheroidal galaxies.

1

Axino dark matter with low reheating temperature

Authors: Krzysztof Turzyski¹; Leszek Roszkowski²; Sebastian Trojanowski²

¹ *University of Warsaw*

² *National Centre for Nuclear Research, Poland*

Corresponding Author: s.trojanowski@ncbj.gov.pl

We examine axino dark matter in the regime of a low reheating temperature T_R after inflation and taking into account that reheating is a non-instantaneous process. This can have a significant effect on the dark matter abundance, mainly due to entropy production in inflaton decays. We study both thermal and non-thermal production of axinos in the context of the MSSM with ten free parameters. We identify the ranges of the axino mass and the reheating temperature allowed by the LHC and other particle physics data in different models of axino interactions. We confront these limits with cosmological constraints coming from the observed dark matter density, large structures formation and big bang nucleosynthesis. We find a number of differences in the phenomenologically acceptable values of the axino mass and the reheating temperature relative to previous studies. In particular, an upper bound on the axino mass becomes dependent on T_R , reaching a maximum value at $T_R \sim 10^2$ GeV. If the lightest ordinary supersymmetric particle is a wino or a higgsino, we obtain lower a limit of approximately 10 GeV for the reheating temperature. We demonstrate also that entropy production during reheating affects the maximum allowed axino mass and lowest values of the reheating temperature.

4

4-jet production in high-energy factorization

Authors: Andreas van Hameren¹; Antoni Szczurek²; Krzysztof Kutak³; Mirko Serino⁴; Rafal Maciula²

¹ *IFJ PAN*

² *IFJ, PAN*

³ *Instytut Fizyki Jądrowej Polskiej Akademii Nauk*

⁴ *Institute of Nuclear Physics, Polish Academy of Sciences*

Corresponding Author: mirko.serino@ifj.edu.pl

Recent progress in the computation of gauge-invariant scattering amplitudes with off-shell particles allows us to perform the first complete High Energy Factorization study of 4-jet production, including all the single and double parton scattering channels. We examine the differences between the two approaches and motivate the importance of using asymmetric cuts in order to pin down multi-parton interactions in future experimental analysis of 4-jet events.

5

Advances in amplitudes with off-shell partons

Author: Andreas van Hameren¹

¹ *IFJ PAN*

Corresponding Author: hameren@ifj.edu.pl

Factorization prescriptions that take into account the transverse momentum of the initial-state partons require hard scattering amplitudes with off-shell initial-state partons. We report on recent advances in the computation of such amplitudes, both at tree-level and at one loop.

6

Clocks and dynamics in Quantum Universe

Author: Przemysław Małkiewicz¹

¹ *APC, Universite Paris Diderot*

Corresponding Author: pmalkiew@gmail.com

Dynamics of general relativistic systems is given with respect to internal clocks. I employ the so called reduced phase space quantization of canonical relativity to show how the choice of the internal clock determines the properties of quantum gravity. For illustration I make use of a semiclassical theory of a FRW model, which replaces the classical singularity with a bounce. As I show, switching the internal clock, on which the quantization is based, can alter the resultant semiclassical theory in almost any way. I conclude by pointing to possible consequences of this result.

7

Probing TeV scale physics with neV neutrons

Author: Kazimierz Bodek¹

¹ *Jagiellonian University in Krakow, Institute of Physics*

Corresponding Author: kazimierz.bodek@uj.edu.pl

The observed excess of matter over antimatter in the Universe remains one of the most important puzzles in physics and unknown sources of fundamental CP symmetry violation are its necessary precondition. Intensive searches for new physics beyond the Standard Model (BSM) concentrate on two main frontiers: (i) high energy experiments performed at collider accelerators and (ii) low energy precision experiments. While the first group looks for exotic particles produced on-shell in high

energy collisions, the second ones seek for tiny deviations in low energy observables which can be attributed to non-existing in the SM exotic interactions. In specific scenarios, low-energy precision experiments offer the insight into the mass scale which is hardly reachable or not reachable at all at present high energy colliders. Slow and very slow neutrons play a prominent role at the low-energy (or precision) frontier providing as many as 20 observables reach in physical information. Electric dipole moment (EDM), decay correlations, n - \bar{n} and n - n' oscillation parameters probe BSM physics. Exceptional properties of precision experiments with cold and ultra-cold neutrons also allow for searching of Lorentz Invariance violation and extra dimensions. Presentation will briefly address recent achievements of fundamental neutron physics.

8

Relaxation rates and phase transitions

Author: Jakub Jankowski¹

¹ *Jagiellonian University*

Corresponding Author: jakubj@th.if.uj.edu.pl

Using a bottom-up gauge gravity constructions, relaxation rates of strongly coupled field theories are computed. A variety of phase structures are considered, from a crossover up to a first order phase transition.

It is established that near the transition the applicability of a hydrodynamic description breaks down at lower momenta than in the conformal case. In the case of the first order phase transition, a spinodal region

appears at temperatures for which the speed of sound squared is negative. An estimate of the preferential scale attained by the unstable modes is also given. Additionally we observe a novel diffusive regime for sound modes over a range of wavelengths. Additionally, for a range of wavelengths we observe a novel diffusive regime for sound modes.

9

Maximum tension principle and entropic force

Author: Hussain Gohar¹

Co-author: Mariusz Dabrowski²

¹ *Institute of Physics, University of Szczecin, Poland*

² *Institute of Physics*

Corresponding Author: hunzaie@gmail.com

We find the series of example theories for which the relativistic limit of maximum tension $F_{\text{max}} = c^4/4G$ represented by the entropic force can be abolished. Among them the varying constants theories, some generalized entropy models applied both for cosmological and black hole horizons as well as some generalized uncertainty principle models.

Summary:

This talk is based on our recently published paper "Abolishing the Maximum tension principle, Phys.Lett. B748 (2015) 428-431"

10

The evolution of cosmological domain walls from Higgs effective potential

Author: Tomasz Krajewski¹

Co-authors: Marek Lewicki¹; Paweł Olszewski¹; Zygmunt Lalak²

¹ *University of Warsaw*

² *Wydział Fizyki UW*

Corresponding Author: tomasz.krajewski@fuw.edu.pl

The study of the renormalization group improved effective potential of the Standard Model has revealed the existence of second minimum at very high field strengths. The existence of the local maximum separating two minima makes the production of cosmological domain walls in the early Universe possible. We investigated the evolution of networks of domain walls using lattice simulations. Our recent results will be presented.

11

Dark matter - how well can one determine its properties?

Author: Leszek Roszkowski¹

Co-authors: Andrew Williams¹; Enrico Maria Sessolo¹; Sebastian Trojanowski¹

¹ *NCBJ*

Corresponding Author: leszekr1@gmail.com

Even if a dark matter signal is measured in direct or indirect search experiments, or both, working out ensuing WIMP properties may prove to be quite challenging. I will discuss some possible outcomes.

12

First detection of a gravitational wave signal

Author: Andrzej Krolak¹

¹ *National Centre for Nuclear Studies*

Corresponding Author: krolak@impan.pl

Detection of a gravitational wave signal from merger of two black holes is one of greatest discoveries of this century. It will open a new window on the Universe. I shall present several aspects of this discovery: gravitational wave detectors, signal modeling and data analysis. I shall mention the follow-up observations of this event by radio, optical, near-infrared, X-ray, and gamma-ray wavelengths with ground- and space-based facilities. I shall describe consequences of this result for physics of fundamental interactions.

13

Recent ATLAS and CMS results on the Standard Model and the Higgs boson

Corresponding Author: artur.kalinowski@fuw.edu.pl

14

Beyond the Standard Model searches in the Higgs sector from ATLAS and CMS

Corresponding Author: pawel.bruckman@ifj.edu.pl

16

Study of matter at extreme conditions with ALICE

Corresponding Author: jacek.otwinowski@ifj.edu.pl

LHC session / 17

Recent ATLAS and CMS results on the Standard Model and the Higgs boson

Corresponding Author: artur.kalinowski@fuw.edu.pl

LHC session / 18

Beyond the Standard Model searches in the Higgs sector from ATLAS and CMS

Corresponding Author: pawel.bruckman@ifj.edu.pl

LHC session / 19

Study of matter at extreme conditions with ALICE

Corresponding Author: jacek.otwinowski@ifj.edu.pl

Cosmology session / 20

First detection of a gravitational wave signal

Corresponding Author: krolak@impan.pl

Cosmology session / 21

Clocks and dynamics in Quantum Universe

Corresponding Author: pmalkiew@gmail.com

Cosmology session / 22

Dark matter - how well can one determine its properties?

Corresponding Author: leszekr1@gmail.com

Cosmology session / 23

Dark matter near a resonance

Corresponding Author: mateusz.duch@fuw.edu.pl

Cosmology session / 24

The averaging problem in cosmology

Cosmology session / 25

Maximum tension principle and entropic force

Corresponding Author: hunzaie@gmail.com

Strong interactions session / 26

Uncovering BFKL dynamics in production of Mueller-Navelet jets at the LHC

Corresponding Author: lech.szymanowski@ncbj.gov.pl

Strong interactions session / 27

Soft gluon resummation in associated Higgs boson production

Strong interactions session / 28

Forward Physics with the ATLAS Detector. Results, Prospects and Polish Contribution

Corresponding Author: rafal.staszewski@ifj.edu.pl

Strong interactions session / 29

From exclusive production, through diffraction, to jets correlations - forward physics results from CMS

Corresponding Author: grzegorz.brona@fuw.edu.pl

Strong interactions session / 30

Status of exotic four-quark mesons

Strong interactions session / 31

Relaxation rates and phase transitions

Corresponding Author: jakubj@th.if.uj.edu.pl

Theory - miscellany / 32

Effective particles in quantum field theory

Corresponding Author: stglazek@fuw.edu.pl

Theory - miscellany / 33

Lattice QCD: a numerical tool for precision physics in the Standard Model and beyond

Corresponding Author: piotr.korcyl@ur.de

Theory - miscellany / 34

Conformal Standard Model

Corresponding Author: meissner@fuw.edu.pl

37

The averaging problem in cosmology

Author: Jan Ostrowski¹

¹ *Uniwersytet Mikołaja Kopernika w Toruniu*

Corresponding Author: ostrowski@astro.uni.torun.pl

Modern cosmology relies strongly on the assumption of isotropy and homogeneity of the Universe. The difficulties arise when this assumption is applied to Einstein equation. This is problematic, since in general, the Einstein tensor built from the averaged metric is not equal to the averaged stress-energy tensor. In this context, the discrepancy between these quantities is called 'the backreaction' and has been a subject of scientific debate among cosmologists and relativists for more than 20 years. In my talk I will present one of the methods to tackle this problem, i.e. the Buchert scalar averaging, together with its applications and possible drawbacks.

38

In search of higher twist effects in proton structure

Author: Leszek Motyka¹

¹ *Jagiellonian University*

Corresponding Author: leszek.motyka@uj.edu.pl

Higher twist effects may provide new, non-trivial insight into proton structure, including multi-parton correlations in the proton, and also into multiple scattering mechanism in QCD. In this talk an overview will be presented of possible higher twist effects in inclusive and diffractive DIS, and in the forward Drell-Yan process.

Beyond the SM session / 39

Hints for the BSM Physics at the LHC

Corresponding Author: stefan.pokorski@fuw.edu.pl

Beyond the SM session / 40

BSM physics at e⁺e⁻ colliders

Corresponding Author: zarnecki@fuw.edu.pl

Beyond the SM session / 41

Phenomenology of Gauge Mediation models at the LHC and future colliders

Corresponding Author: tomasz.jelinski@us.edu.pl

Beyond the SM session / 42

Enabling Electroweak Baryogenesis through Dark Matter

Corresponding Author: marek.lewicki@fuw.edu.pl

Radiative corrections at colliders session / 45

Successes and challenges in perturbative QCD

Corresponding Author: mczaikon@physik.rwth-aachen.de

Radiative corrections at colliders session / 46

Matching fixed order QCD with parton shower for Drell-Yan and Higgs production

Radiative corrections at colliders session / 47

Advances in amplitudes with off-shell partons

Corresponding Author: hameren@ifj.edu.pl

Radiative corrections at colliders session / 48

Radiative corrections to hadron production in e^+e^- annihilation

Corresponding Author: szymon885@gazeta.pl

49

4-jet production in high-energy factorization

Corresponding Author: mirko.serino@ifj.edu.pl

Radiative corrections at colliders session / 50

4-jet production in high-energy factorization

Corresponding Author: mirko.serino@ifj.edu.pl

Quark-flavour physics session / 51

Probing TeV scale physics with neV neutrons

Corresponding Author: kazimierz.bodek@uj.edu.pl

Quark-flavour physics session / 52

Recent Belle results

Corresponding Author: yjkwon63@yonsei.ac.kr

Quark-flavour physics session / 53

Recent results from LHCb

Corresponding Author: mchrzasz@cern.ch

54

Recent results from LHCb

Corresponding Author: mchrzasz@cern.ch

Quark-flavour physics session / 55

Belle II perspective

Quark-flavour physics session / 56

LHCb perspectives on flavour physics

Corresponding Author: amucha@agh.edu.pl

Neutrino Physics session / 57

Neutrino properties determined in oscillation experiments

Corresponding Author: ewa.rondio@ncbj.gov.pl

Neutrino Physics session / 58

Astrophysical searches with neutrino detectors

Neutrino Physics session / 59

Neutrino portal to new physics

Corresponding Author: bhupaldev@gmail.com