

ENERGY SPECTRUM OF MAGNETIC FIELDS IN THE Universe at the EWPT

Department for Theoretical Physics





Table of Contents

- 1. Introduction to primordial magnetic fields
- 2. Magnetic fields from the electroweak phase transition
- 3. Lattice simulations vs Analytic approach
- 4. Continuous case and the form of spectra

Introduction

- Galaxies have observable magnetic fields of $B_{\rm galaxy} \sim \mathcal{O}(1-10)~\mu{
 m G}$ [Unger et al., 2024]
 - $\,\triangleright\,$ Amplification of weak "seed magnetic fields" by the rotating galactic plasma
 - ▶ But what is the origin of this seed field?



INTRODUCTION

- Galaxies have observable magnetic fields of $B_{\rm galaxy} \sim \mathcal{O}(1-10) \,\mu G$ [Unger et al., 2024]
 - > Amplification of weak "seed magnetic fields" by the rotating galactic plasma
 - ▶ But what is the origin of this seed field?
- Evidence for weak galactic fields outside of galaxies (in voids)
 - Observed in TeV Blazar data [Neronov et al., 2010]
 - Diffuse image caused by magnetic fields between the blazar and observer
 - \triangleright Result: $B_{\text{void}} \sim \mathcal{O}(10^{-16})$ G field with $\lambda_{\text{void}} \sim 1$ Mpc correlation length
 - \triangleright Compare: $\lambda_{\rm void} \sim 10^{21} \,\mathrm{m} \longrightarrow \text{Weak field but enormous distances}$



INTRODUCTION

- Galaxies have observable magnetic fields of $B_{\rm galaxy} \sim \mathcal{O}(1-10) \,\mu G$ [Unger et al., 2024]
 - > Amplification of weak "seed magnetic fields" by the rotating galactic plasma
 - ▶ But what is the origin of this seed field?
- Evidence for weak galactic fields outside of galaxies (in voids)
 - Described in TeV Blazar data [Neronov et al., 2010]
 - Diffuse image caused by magnetic fields between the blazar and observer
 - \triangleright Result: $B_{\text{void}} \sim \mathcal{O}(10^{-16})$ G field with $\lambda_{\text{void}} \sim 1$ Mpc correlation length
 - \triangleright Compare: $\lambda_{\text{void}} \sim 10^{21} \,\text{m} \longrightarrow \text{Weak field but enormous distances}$
- Weak and homogeneous fields → Possibly early Universe origin
 - ▶ A possible explanation is through topological effects [Vachaspati, 1991]
 - Despite the absence of stable defects in the SM, unstable structures-called Nambu dumbbells-may exist [Nambu, 1977]

PRIMORDIAL MAGNETIC FIELDS

$$\mathcal{L}_{\text{SM}} = |D_{\mu}\Phi|^{2} + V(\Phi) + \dots$$

$$\Phi_{\text{min}}: D_{\mu}\Phi_{\text{min}} \equiv \left(\partial_{\mu} - \frac{\mathrm{i}g}{2}\tau^{a}A_{\mu}^{a} - \frac{\mathrm{i}g'}{2}A_{\mu}^{0}\right)\Phi_{\text{min}} = 0 \quad \text{and} \quad V(\Phi_{\text{min}}) = 0.$$

$$g'A_{\mu}^{0} = -\mathrm{i}\sin^{2}\theta_{w}\left(\Phi^{\dagger}\cdot\partial_{\mu}\Phi - \partial_{\mu}\Phi^{\dagger}\cdot\Phi\right)$$

$$gA_{\mu}^{a} = -\mathrm{i}\cos^{2}\theta_{w}n^{a}\left(\Phi^{\dagger}\cdot\partial_{\mu}\Phi - \partial_{\mu}\Phi^{\dagger}\cdot\Phi\right) - \epsilon^{abc}n^{b}\partial_{\mu}n^{c} \qquad \left(n^{a} = \Phi^{\dagger}\cdot\tau^{a}\cdot\Phi\right)$$

$$\mathcal{F}_{\mu\nu} = \cos\theta_{w}F_{\mu\nu}^{0} - \sin\theta_{w}F_{\mu\nu}^{a}n^{a} = -\frac{2\mathrm{i}}{g}\sin\theta_{w}\left(\partial_{\mu}\Phi^{\dagger}\partial_{\nu}\Phi - \partial_{\nu}\Phi^{\dagger}\partial_{\mu}\Phi\right).$$

$$\mathbf{B} = -\mathrm{i}\frac{2\sin\theta_{w}}{a}\left(\nabla\Phi^{\dagger}\times\nabla\Phi\right) \quad \text{and} \quad \mathbf{A} = -\mathrm{i}\frac{2\sin\theta_{w}}{a}\Phi^{\dagger}\nabla\Phi$$

CORRELATION FUNCTIONS

- \bullet To describe the statistical features of the $\mathbf{B}(\mathbf{r})$ field we use **correlation functions**
 - \triangleright Assuming isotropy and $\nabla \cdot \mathbf{B} = 0$ (defects have decayed)
 - ▶ Spatial correlation function:

$$\langle B_i(\mathbf{x} + \mathbf{r})B_j(\mathbf{r})\rangle = M_{\rm N}(r)P_{ij} + M_{\rm L}(r)\hat{r}_i\hat{r}_j + \epsilon_{ijk}\hat{r}_kM_{\rm H}(r)$$

▶ Fourier transform convention:

$$\mathbf{b}(\mathbf{k}) = \int \mathrm{d}^3 r \, \mathbf{B}(\mathbf{r}) \exp\left(\mathrm{i} \mathbf{k} \cdot \mathbf{r}\right)$$

CORRELATION FUNCTIONS

- To describe the statistical features of the $\mathbf{B}(\mathbf{r})$ field we use **correlation functions**
 - \triangleright Assuming isotropy and $\nabla \cdot \mathbf{B} = 0$ (defects have decayed)
 - ▶ Spatial correlation function:

$$\langle B_i(\mathbf{x} + \mathbf{r})B_j(\mathbf{r})\rangle = M_{\rm N}(r)P_{ij} + M_{\rm L}(r)\hat{r}_i\hat{r}_j + \epsilon_{ijk}\hat{r}_kM_{\rm H}(r)$$

▶ Fourier-space correlation function:

$$\langle b_i(\mathbf{k})b_j(\mathbf{k}')\rangle = (2\pi)^6 \delta^{(3)}(\mathbf{k} - \mathbf{k}') \left[\frac{E_{\mathbf{M}}(k)}{4\pi k^2} \mathbf{P}_{ij}^{(k)} + \mathrm{i}\epsilon_{ijl}k_l \frac{H_{\mathbf{M}}(k)}{8\pi k^2} \right]$$

CORRELATION FUNCTIONS

- \bullet To describe the statistical features of the $\mathbf{B}(\mathbf{r})$ field we use **correlation functions**
 - \triangleright Assuming isotropy and $\nabla \cdot \mathbf{B} = 0$ (defects have decayed)
 - ▶ Spatial correlation function:

$$\langle B_i(\mathbf{x} + \mathbf{r})B_j(\mathbf{r})\rangle = M_{\rm N}(r)P_{ij} + M_{\rm L}(r)\hat{r}_i\hat{r}_j + \epsilon_{ijk}\hat{r}_kM_{\rm H}(r)$$

▶ Fourier-space correlation function:

$$\langle b_i(\mathbf{k})b_j(\mathbf{k}')\rangle = (2\pi)^6 \delta^{(3)}(\mathbf{k} - \mathbf{k}') \left[\frac{E_{\mathbf{M}}(k)}{4\pi k^2} \mathbf{P}_{ij}^{(k)} + \mathrm{i}\epsilon_{ijl}k_l \frac{H_{\mathbf{M}}(k)}{8\pi k^2} \right]$$

 \triangleright Trace of these \rightarrow correlation function and power spectrum



Energy spectrum

▶ Using the ergodic principle we define the **correlation function**:

$$\xi(\mathbf{r}) \equiv \langle \mathbf{B}(\mathbf{r}) \cdot \mathbf{B}(\mathbf{x} + \mathbf{r}) \rangle = \frac{1}{V} \int_{V} d^{3}x \, \mathbf{B}(\mathbf{x}) \cdot \mathbf{B}(\mathbf{x} + \mathbf{r})$$

▶ Using $V = (2\pi)^3 \delta^{(3)}(\mathbf{k} = \mathbf{0})$ we extract the **energy spectrum**:

$$E_{\mathbf{M}}(\mathbf{k}) = \frac{k^2 \langle |\mathbf{b}(\mathbf{k})|^2 \rangle}{(2\pi)^2 V}$$

▶ Relationship between the correlation function and the energy spectrum:

$$\xi(\mathbf{r}) = \frac{1}{2} \int_0^\infty \mathrm{d}k \, E_\mathrm{M}(k) j_0(kr) \quad \longleftrightarrow \quad E_\mathrm{M}(k) = \frac{k^2}{\pi} \int_0^\infty \mathrm{d}r \, r^2 \xi(\mathbf{r}) j_0(kr)$$

Magnetic fields from EWPT

EWPT AND THE SCALAR FIELD

• After EWPT the scalar field $\Phi(\mathbf{r}) \equiv v\tilde{\Phi}(\mathbf{r})$ is on its vacuum manifold:

$$|\tilde{\Phi}|^2 = \tilde{\phi}_1^2 + \tilde{\phi}_2^2 + \tilde{\phi}_3^2 + \tilde{\phi}_4^2 \stackrel{!}{=} 1 \longrightarrow \tilde{\Phi} \sim S_3$$

• Side note: topological defects require non-trivial homotopy classes, however

$$\pi_0(S_3) = \pi_1(S_3) = \pi_2(S_3) = 0 \longrightarrow \text{No topological defects!}$$

• Hopf-parametrization of the scalar field on S_3 :

$$\tilde{\Phi} = \begin{pmatrix} \cos \alpha \exp(\mathrm{i}\beta) \\ \sin \alpha \exp(\mathrm{i}\gamma) \end{pmatrix}, \quad \text{where} \quad \beta, \gamma \in [0, 2\pi] \quad \text{and} \quad 2u \equiv \cos(2\alpha) \in [-1, 1]$$

• Hopf-parameters u, β, γ distributed uniformly at random

LATTICE SETUP I.

- Idea based on [Vachaspati et al., 2025]
- Cubic lattice of size N^3 with lattice spacing δx and $L = N\delta x = V^{1/3}$
- Choose random Hopf-angles α, β, γ at each lattice site:

 $\tilde{\Phi}(\mathbf{x}_{ijk})$, where $\mathbf{x}_{ijk} = i\hat{\mathbf{x}} + j\hat{\mathbf{y}} + k\hat{\mathbf{z}}$ Random realization of the scalar field:

- Let $2\sin\theta_{\rm w}/q \to 1$ so that $\mathbf{A} = -\mathrm{i}\tilde{\Phi}^{\dagger}\nabla\tilde{\Phi}$
- Define the vector potential through the link variable:

$$T_a(\mathbf{x}) \equiv \mathbf{A}(\mathbf{x}) \cdot (\delta x \, \hat{\mathbf{a}}) \simeq \mathfrak{Im} \big[\tilde{\Phi}^{\dagger}(\mathbf{x}) \cdot \tilde{\Phi}(\mathbf{x} + \delta x \, \hat{\mathbf{a}}) \big] \equiv \mathbf{x} \qquad \mathbf{x} + \delta x \, \hat{\mathbf{a}}$$

Important step: **Periodic boundary condition** when defining link variables!



LATTICE SETUP II.

Magnetic field defined via the loop integral:


$$\mathbf{B}(\mathbf{x}^p)\delta x^2 = \hat{\mathbf{p}} \oint_{\partial p} \mathrm{d}\mathbf{l} \cdot \mathbf{A}(\mathbf{x})$$

On the lattice this reduces to a sum of 4 link variables, e.g.:


$$B_{x}(\mathbf{x}_{i,j,k}^{p})\delta x^{2} = \mathbf{x}_{i,j,k}^{p}$$

$$= T_{y}(\mathbf{x}_{i,j,k}) + T_{z}(\mathbf{x}_{i,j+1,k}) + T_{-y}(\mathbf{x}_{i,j+1,k+1}) + T_{-z}(\mathbf{x}_{i,j,k+1})$$

VISUALIZING THE LATTICE



BASIC SIMULATION APPROACH



"SIMULATION ON PAPER"

- Basic simulation approach: Slow for large lattice size N and very noisy for small k (large distances)
 - \triangleright Personal problem: cannot run on a laptop due to limited RAM for $N \ge 256$

"SIMULATION ON PAPER"

- Basic simulation approach: Slow for large lattice size N and very noisy for small k (large distances)
 - \triangleright Personal problem: cannot run on a laptop due to limited RAM for $N \gtrsim 256$
- We can do much better!
 - ▶ We can exploit the randomness of $\Phi(\mathbf{x}_{ijk})$ and the law of large numbers (nr. of lattice points $\gg 1$)
 - ▶ We can also use general results in the mathematics of Fourier transform

"SIMULATION ON PAPER"

- Basic simulation approach: Slow for large lattice size N and very noisy for small k (large distances)
 - \triangleright Personal problem: cannot run on a laptop due to limited RAM for $N \gtrsim 256$

• We can do much better!

- ▶ We can exploit the randomness of $\Phi(\mathbf{x}_{ijk})$ and the law of large numbers (nr. of lattice points $\gg 1$)
- ▶ We can also use general results in the mathematics of Fourier transform

• Approach:

- 1. Find the spatial correlation function \rightarrow
- 2. Average over polar angles to get the isotropic correlation function \rightarrow
- 3. Use relationship between power spectrum and correlation function \rightarrow

 $\xi({f r})$

 $\xi_{\rm iso}(r)$

 $E_{\rm M}(k)$

ANALYTIC APPROACH FOR THE SPECTRUM

- Approach:
 - \triangleright Evaluate the correlation function $\xi(\mathbf{r})$ for the lattice:

$$\xi(\mathbf{r}) = \frac{1}{N^3} \sum_{i=1}^3 \sum_{\mathbf{x}} B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \simeq \sum_{i=1}^3 \langle B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \rangle_{\mathbf{x}}$$

ightharpoonupUsing $\langle T_a(\mathbf{x})T_b(\mathbf{y})\rangle = 0$ unless $T_a(\mathbf{x})$ and $T_b(\mathbf{y})$ are the "same" link:

$$\langle B_x(\mathbf{x}_{i,j-1,k}^p) B_x(\mathbf{x}_{i,j,k}^p) \rangle = \langle (\mathbf{i},\mathbf{j},\mathbf{i},\mathbf{k}+1) & (\mathbf{i},\mathbf{j},\mathbf{k}+1) & (\mathbf{i},\mathbf{j}+1,\mathbf{k}+1) \rangle \\ = \langle (\mathbf{i},\mathbf{j},\mathbf{k}) & (\mathbf{i},\mathbf{j},\mathbf{k}+1) \rangle \\ = \langle T_z(\mathbf{x}_{i,j,k}) T_{-z}(\mathbf{x}_{i,j,k+1}) \rangle \equiv -\langle [T_z(\mathbf{x}_{i,j,k})]^2 \rangle$$



Analytic Approach – The Correlation Function

- Approach:
 - Evaluate the correlation function $\xi(\mathbf{r})$ for the lattice:

$$\xi(\mathbf{r}) = \frac{1}{N^3} \sum_{i=1}^3 \sum_{\mathbf{x}} B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \simeq \sum_{i=1}^3 \langle B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \rangle_{\mathbf{x}}$$

 \triangleright Using $\langle T_a(\mathbf{x})T_b(\mathbf{y})\rangle = 0$ unless $T_a(\mathbf{x})$ and $T_b(\mathbf{y})$ are the same link, e.g.:

$$\langle B_x(\mathbf{x}_{i,j-1,k}^p) B_x(\mathbf{x}_{i,j,k}^p) \rangle = -\langle [T_z(\mathbf{x}_{i,j,k})]^2 \rangle = -\frac{1}{4}$$

ANALYTIC APPROACH – THE CORRELATION FUNCTION

- Approach:
 - Evaluate the correlation function $\xi(\mathbf{r})$ for the lattice:

$$\xi(\mathbf{r}) = \frac{1}{N^3} \sum_{i=1}^3 \sum_{\mathbf{x}} B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \simeq \sum_{i=1}^3 \langle B_i(\mathbf{x}) B_i(\mathbf{x} + \mathbf{r}) \rangle_{\mathbf{x}}$$

 \triangleright Using $\langle T_a(\mathbf{x})T_b(\mathbf{y})\rangle = 0$ unless $T_a(\mathbf{x})$ and $T_b(\mathbf{y})$ are the same link, e.g.:

$$\langle B_x(\mathbf{x}_{i,j-1,k}^p) B_x(\mathbf{x}_{i,j,k}^p) \rangle = -\langle [T_z(\mathbf{x}_{i,j,k})]^2 \rangle = -\frac{1}{4}$$

Obviously, correlations vanish for $r > \delta x$ as there will be no shared links:

$$\xi(\mathbf{r}) = \frac{1}{\delta x^4} \sum_{i=1}^{3} \left(\delta_{\mathbf{r},\mathbf{0}} - \frac{1}{4} \sum_{j=1}^{3} \delta_{\mathbf{r},\pm \delta x \hat{\mathbf{e}}_j} (1 - \delta_{ij}) \right)$$



DISCRETE APPROACH I.

- Isotropic limit on the lattice:
 - Introduce the set of lattice points that are at a given distance:

$$\mathfrak{P}(r) = \{ \mathbf{r} \mid |\mathbf{r}| = r; \, \mathbf{r} \in \text{lattice} \} \qquad \rightarrow \qquad |\mathfrak{P}(0)| = 1, \, |\mathfrak{P}(\delta x)| = 6$$

 \triangleright We average over the direction-dependence of $\xi(\mathbf{r})$:

$$\xi_{\rm iso}(r) = \frac{3}{\delta x^4 |\mathcal{P}(r)|} \left(\delta_{r,0} - \delta_{r,\delta x} \right) = \frac{3}{\delta x^4} \left(\delta_{r,0} - \frac{1}{6} \delta_{r,\delta x} \right)$$

The power spectrum is the DFT of the correlation function:

$$\langle |\mathbf{b}(\mathbf{k})|^2 \rangle = V \delta x^3 \sum_{\mathbf{r}} \xi(\mathbf{r}) \exp(i\mathbf{k} \cdot \mathbf{r}) = \delta x^2 N^3 \sum_{i=1}^3 \left[1 - \cos\left(k_i \delta x\right) \right]$$



DISCRETE APPROACH I.

- Isotropic limit on the lattice:
 - Introduce the set of lattice points that are at a given distance:

$$\mathcal{P}(r) = \{\mathbf{r} \mid |\mathbf{r}| = r; \mathbf{r} \in \text{lattice}\}$$
 \rightarrow $|\mathcal{P}(0)| = 1, |\mathcal{P}(\delta x)| = 6$

We neglect the direction-dependence of the 2nd term in $\xi(\mathbf{r})$:

$$\xi_{\rm iso}(r) = \frac{3}{\delta x^4 |\mathcal{P}(r)|} \left(\delta_{r,0} - \delta_{r,\delta x} \right) = \frac{3}{\delta x^4} \left(\delta_{r,0} - \frac{1}{6} \delta_{r,\delta x} \right)$$

The power spectrum is the DFT of the correlation function:

$$\langle |\mathbf{b}(\mathbf{k})|^2 \rangle = \delta x^2 N^3 \sum_{i=1}^{3} \left[1 - \left(1 - \frac{k_i^2 L^2}{2N^2} + \mathcal{O}(k_i^4 L^4) \right) \right] = \frac{N}{2} \delta x^2 (kL)^2 \left[1 + \mathcal{O}\left((kL)^2 \right) \right]$$

DISCRETE APPROACH II.

• The energy spectrum is obtained from the power spectrum:


$$8\pi^2 \frac{E_{\rm M}(k)}{k^4} = \delta x \left[1 + \mathcal{O}\left((kL)^2\right)\right] \qquad \to \qquad E_{\rm M}(k) \propto k^4$$

DISCRETE APPROACH II.

• The energy spectrum is obtained from the power spectrum:

$$8\pi^2 \frac{E_{\rm M}(k)}{k^4} = \delta x \left[1 + \mathcal{O}\left((kL)^2\right) \right] \qquad \to \qquad E_{\rm M}(k) \propto k^4$$

- ullet More detailed spectrum \to generalized coarse-grain algorithms
 - \triangleright Effectively: averaging over $n \times n$ sized plaquettes
 - \triangleright Correlations will exist up to larger distances for n^2 -plaquette averaged fields, e.g.:



DISCRETE APPROACH II.

• The energy spectrum is obtained from the power spectrum:

$$8\pi^2 \frac{E_{\mathcal{M}}(k)}{k^4} = \delta x \left[1 + \mathcal{O}\left((kL)^2\right) \right] \qquad \to \qquad E_{\mathcal{M}}(k) \propto k^4$$

- More detailed spectrum \rightarrow generalized coarse-grain algorithms
 - \triangleright Effectively: averaging over $n \times n$ sized plaquettes
 - \triangleright Correlations will exist up to larger distances for n^2 -plaquette averaged fields
 - ▶ More complicated correlation function and power spectrum...

$$\xi_{\text{iso}}^{(4\text{PA})}(r) = \frac{2 \cdot 3}{(2\delta x)^4 |\mathcal{P}(r)|} \left[\delta_{r,0} + \delta_{r,\delta x} - \delta_{r,2\delta x} - \delta_{r,\sqrt{5}\delta x} \right]$$

 \triangleright Scaling of $E_{\rm M}(k)$ remains the same for small k



CONTINUOUS APPROACH

• Starting point is the same $\xi(\mathbf{r})$ lattice correlation function:

$$\xi_{\rm iso}(r) = \frac{3}{\delta x^4 |\mathcal{P}(r)|} (\delta_{r,0} - \delta_{r,\delta x})$$

- \triangleright Lattice: $|\mathcal{P}(r)|$ counts nr. of neighbors
- \triangleright Lattice: $\delta_{r,a}$ Kronecker-delta

Continuous Approach

• Starting point is the same $\xi(\mathbf{r})$ lattice correlation function:

$$\xi_{\rm iso}(r) = \frac{3}{\delta x^4 |\mathcal{P}(r)|} \left(\delta_{r,0} - \delta_{r,\delta x} \right) \qquad \rightarrow \qquad \left[\frac{\xi_{\rm iso}^{(c)}(r)}{4\pi r^2 \delta x} \left(\delta(r) - \delta(r - \delta x) \right) \right]$$

- ▶ Lattice: $|\mathcal{P}(r)|$ counts nr. of neighbors \rightarrow Continuum: $|\mathcal{P}(r)| \rightarrow 4\pi r^2/\delta x^2$
- \triangleright Lattice: $\delta_{r,a}$ Kronecker-delta \rightarrow Continuum: $\delta_{r,a} \rightarrow \delta x \, \delta(r-a)$

Continuous Approach

• Starting point is the same $\xi(\mathbf{r})$ lattice correlation function:

$$\xi_{\rm iso}(r) = \frac{3}{\delta x^4 |\mathcal{P}(r)|} \left(\delta_{r,0} - \delta_{r,\delta x} \right) \qquad \rightarrow \qquad \left[\frac{\xi_{\rm iso}^{(c)}(r)}{4\pi r^2 \delta x} \left(\delta(r) - \delta(r - \delta x) \right) \right]$$


- ▶ Lattice: $|\mathcal{P}(r)|$ counts nr. of neighbors \rightarrow Continuum: $|\mathcal{P}(r)| \rightarrow 4\pi r^2/\delta x^2$
- ightharpoonup Lattice: $\delta_{r,a}$ Kronecker-delta ightharpoonup Continuum: $\delta_{r,a}
 ightharpoonup \delta x \, \delta(r-a)$
- The energy spectrum is the Hänkel transform of $\xi_{\text{ico}}^{(c)}(r)$:

$$E_{\rm M}(k) = \frac{k^2}{\pi} \int_{-\pi}^{\infty} dr \, r^2 \xi_{\rm iso}^{(c)}(r) j_0(kr)$$

$$8\pi^2 \frac{E_{\rm M}(k)}{k^4} = \frac{6}{k^2 \delta x} \left(1 - j_0(k \delta x) \right) \quad \to \quad E_{\rm M}(k) \propto k^4$$



SIMULATION VS ANALYTIC SOLUTION




Going further: Interpolated Fields

- There remain some issues with the above lattice approach:
 - ▶ Crudeness of derivatives (link variables)
 - ▶ Insensitivity for scales below the lattice spacing
 - ▶ Regularity of the cubic lattice

GOING FURTHER: INTERPOLATED FIELDS

- There remain some issues with the above lattice approach:
 - ▶ Crudeness of derivatives (link variables)
 - ▶ Insensitivity for scales below the lattice spacing
 - ▶ Regularity of the cubic lattice
- We attempt to solve all of these with the following changes:
 - ▶ Irregular lattice setup
 - ▷ Continuous interpolation of the lattice field
 - → Using "radial basis function (RBF)" interpolation due to irregularity
 - \triangleright Sampling of the continuous field on a dense grid \rightarrow DFT as before...
 - → Technically, direct FT is possible but very slow

EXAMPLE SLICE OF THE CONTINUOUS MAGNETIC FIELD



CAUSAL CORRELATION FUNCTIONS

- Due to the interpolation \longrightarrow nearby lattice sites are not independent
- Need "reasonable" analytic formulae to fit for the correlation function and the power spectrum
- Important detail: Causal correlation functions [Durrer et al., 2003]
 - \triangleright Mathematics: Provided that $\xi(r) = 0$ for $r > \ell$ then $E_{\rm M}(k) \propto k^4$ for small k (actually, exponentially vanishing is enough)
 - Dur above analytic calculation confirmed this explicitly
 - \triangleright If $\xi(r)$ is exponentially small for large r then so is $E_{\rm M}(k)$ for large k
 - Empirical formula for the k^4 energy spectrum with an exponential cutoff:

$$\mathcal{E}(k; \{b_a\}, \sigma) \propto k^4 \left(1 + \sum_{a=1}^{n/2} b_a k^{2a}\right) \exp\left(-\frac{2k^2}{\sigma^2}\right)$$



• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

Energy spectrum \rightarrow Hänkel transform of C(r):

[Fubini's theorem]

$$\mathcal{E}(k) = \frac{1}{\pi} \sum_{n=0}^{n_{\text{max}}} p_n k^{-(n+1)} \int_0^\infty dq \, q^{n+2} \exp(-aq^2/k^2) j_0(q)$$

• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

• Energy spectrum \rightarrow Hänkel transform of C(r):

$$\mathcal{E}(k) = \frac{1}{2\pi} \sum_{n=0}^{\infty} p_n \frac{k^2}{a^{m_n}} \Gamma(m_n) \, {}_{1}F_1\left(m_n; m_0; -\frac{k^2}{4a}\right), \quad \text{where } m_n = \frac{3+n}{2}$$




• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

Energy spectrum \rightarrow Hänkel transform of C(r) with $m_n = (3+n)/2$:

$$\mathcal{E}(k) = \frac{1}{2\pi} \sum_{n=0}^{n_{\text{max}}} p_n \frac{\Gamma(m_n)}{a^{m_n}} \sum_{j=0}^{n/2} \frac{(-1)^j}{(2j+1)!!} \frac{k^{2(j+1)}}{(2a)^j} \binom{n/2}{j} \exp\left(-\frac{k^2}{4a}\right)$$



• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

Energy spectrum \rightarrow Hänkel transform of C(r) with $m_n = (3+n)/2$:

$$\mathcal{E}(k) = \frac{1}{2\pi} \sum_{n=0}^{n_{\text{max}}} p_n \frac{\Gamma(m_n)}{a^{m_n}} \sum_{j=0}^{n/2} \frac{(-1)^j}{(2j+1)!!} \frac{k^{2(j+1)}}{(2a)^j} \binom{n/2}{j} \exp\left(-\frac{k^2}{4a}\right)$$

Compare with the empirical formula:

$$\mathcal{E}(k) \propto k^4 \left(1 + \sum_{i=1}^{n/2} b_a k^{2a}\right) \exp\left(-\frac{2k^2}{\sigma^2}\right)$$

• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

• Energy spectrum \rightarrow Hänkel transform of C(r) with $m_n = (3+n)/2$:

$$\mathcal{E}(k) = \frac{1}{2\pi} \left| \sum_{n=0}^{n_{\text{max}}} p_n \frac{\Gamma(m_n)}{a^{m_n}} \sum_{j=0}^{n/2} \frac{(-1)^j}{(2j+1)!!} \binom{n/2}{j} \frac{k^{2(j+1)}}{(2a)^j} \right| \left[\exp\left(-\frac{k^2}{4a}\right) \right]$$

• Compare with the empirical formula:

$$\mathcal{E}(k) \propto k^4 \left(1 + \sum_{a=1}^{n/2} b_a k^{2a}\right) \left[\exp\left(-\frac{2k^2}{\sigma^2}\right)\right]$$

• What specific form should a causal correlation function take to give $\mathcal{E}(k) \propto k^4$?

$$C(r) = \sum_{n=0}^{n_{\text{max}}} p_n r^n \exp(-ar^2), \text{ with } p_{\text{odd}} = 0$$

Energy spectrum \rightarrow Hänkel transform of C(r) with $m_n = (3+n)/2$:


$$\mathcal{E}(k) = \frac{1}{2\pi} \sum_{n=0}^{\infty} p_n \frac{\Gamma(m_n)}{a^{m_n}} \sum_{j=0}^{n/2} \frac{(-1)^j}{(2j+1)!!} \frac{k^{2(j+1)}}{(2a)^j} \binom{n/2}{j} \exp\left(-\frac{k^2}{4a}\right)$$

• The k^2 term \leftrightarrow the j=0 part of the expression has to cancel:

$$\sum_{n=0}^{n_{\text{max}}} p_n a^{-n/2} \Gamma\left(\frac{3+n}{2}\right) = 0$$

SPECTRUM OF THE PRIMORDIAL MAGNETIC FIELD

• The spatial correlation function and the energy spectrum (normalized to the spectral peak at k_*):





CONCLUSIONS AND OUTLOOK

• Conclusions:

- > We investigated the spectrum of the magnetic field that could be produced around the time of the EWPT
- > We showed that the spectrum may be directly derived without the need of numerical simulations
- \triangleright We confirmed that the spectrum is "causal" and that it scales as $E_{\rm M} \propto k^4$
- ▶ We derived a general formula for the isotropic spectra
- ▶ We showed that the general formula fits the results of the "continuous" simulations

Outlook:

- > The spectrum is evolved using MHD equations to compare to the magnetic field strength of $\mathcal{O}(10^{-16})$ G at $\lambda = 1$ Mpc scales
- > This simulation does not generate helical fields thus the obtained fields today are too weak
- Introduce parity violation initially and get a non-zero helicity spectrum