

Renormalons in heavy quark physics and lattice: the pole mass and the gluon condensate

Antonio Pineda

Universitat Autònoma de Barcelona & IFAE

Matter to the deepest,
Podlesice (Poland), 3-8 September, 2017

Renormalons

Originally (Lautrup, 't Hooft).

Renormalon: summation of "bubbles". Running of α_s .




Figure : Sum of the bubbles in the quark propagator.

Pole mass (Bigi, Shifman, Uraltsev, Vainshtein; Beneke, Braun)

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \quad B_n \sim n!$$

$$\delta m \propto \int dk \alpha(k) \sim \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n \int dk \ln^n \frac{\mu}{k} = \mu \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n n!$$

$$\beta_0^{\text{QED}} = -\frac{4}{3} T_F n_f \rightarrow \beta_0^{\text{QCD}} = \frac{11}{3} C_A - \frac{4}{3} T_F n_f \text{ naive non-abelianization.}$$

Beyond bubbles: renormalization group methods (Parisi; Beneke, ...)

NP OPE (Novikov, Shifman, Vainshtein, Zakharov)

Renormalons

Originally (Lautrup, 't Hooft).

Renormalon: summation of "bubbles". Running of α_s .




Figure : Sum of the bubbles in the quark propagator.

Pole mass (Bigi, Shifman, Uraltsev, Vainshtein; Beneke, Braun)

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \quad B_n \sim n!$$

$$\delta m \propto \int d\mu \alpha(\mu) \sim \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n \int d\mu \ln^n \frac{\mu}{k} = \mu \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n n!$$

$$\beta_0^{\text{QED}} = -\frac{4}{3} T_F n_f \rightarrow \beta_0^{\text{QCD}} = \frac{11}{3} C_A - \frac{4}{3} T_F n_f \text{ naive non-abelianization.}$$

Beyond bubbles: renormalization group methods (Parisi; Beneke, ...)

NP OPE (Novikov, Shifman, Vainshtein, Zakharov)

Renormalons

Originally (Lautrup, 't Hooft).

Renormalon: summation of "bubbles". Running of α_s .




Figure : Sum of the bubbles in the quark propagator.

Pole mass (Bigi, Shifman, Uraltsev, Vainshtein; Beneke, Braun)

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \quad B_n \sim n!$$

$$\delta m \propto \int dk \alpha(k) \sim \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n \int dk \ln^n \frac{\mu}{k} = \mu \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n n!$$

$$\beta_0^{\text{QED}} = -\frac{4}{3} T_F n_l \rightarrow \beta_0^{\text{QCD}} = \frac{11}{3} C_A - \frac{4}{3} T_F n_l \text{ naive non-abelianization.}$$

Beyond bubbles: renormalization group methods (Parisi; Beneke, ...)

NP OPE (Novikov, Shifman, Vainshtein, Zakharov)

Renormalons

Originally (Lautrup, 't Hooft).

Renormalon: summation of "bubbles". Running of α_s .




Figure : Sum of the bubbles in the quark propagator.

Pole mass (Bigi, Shifman, Uraltsev, Vainshtein; Beneke, Braun)

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \quad B_n \sim n!$$

$$\delta m \propto \int dk \alpha(k) \sim \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n \int dk \ln^n \frac{\mu}{k} = \mu \alpha_s(\mu) \sum_{n=0}^{\infty} \left(\frac{\beta_0 \alpha(\mu)}{2\pi} \right)^n n!$$

$$\beta_0^{\text{QED}} = -\frac{4}{3} T_F n_l \rightarrow \beta_0^{\text{QCD}} = \frac{11}{3} C_A - \frac{4}{3} T_F n_l \text{ naive non-abelianization.}$$

Beyond bubbles: renormalization group methods (Parisi; Beneke, ...)

NP OPE (Novikov, Shifman, Vainshtein, Zakharov)

Modern view: EFT/Factorization

$$\mathcal{L} = \sum_n \frac{1}{m^n} c_n O_n \quad c(\nu) = \bar{c} + \sum_{n=0}^{\infty} c_n \alpha_s^{n+1}.$$

The Wilson coefficients are believed to be asymptotic: $c_n \sim n!$

If so such behavior should comply with the Operator Product Expansion.

EFT/factorization definition of renormalon: Asymptotic behavior of the perturbative expansion such that the associated ambiguity in the summation of the perturbative series can be absorbed into a higher order operator.

Example:

$$M_B = m_{\text{OS}} + \bar{\Lambda}_B + \mathcal{O}(1/m_{\text{OS}})$$

M_B is renormalon free. Therefore m_{OS} suffers from renormalon ambiguities:

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots)$$

with $B_n \sim n!$. In other words

$$\delta_{np}^{(\text{pert.})} m_{\text{OS}} = \delta_{np}^{(\text{pert.})} m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \sim \Lambda_{\text{QCD}}!$$

Modern view: EFT/Factorization

$$\mathcal{L} = \sum_n \frac{1}{m^n} c_n O_n \quad c(\nu) = \bar{c} + \sum_{n=0}^{\infty} c_n \alpha_s^{n+1}.$$

The Wilson coefficients are believed to be asymptotic: $c_n \sim n!$

IF SO such behavior should comply with the Operator Product Expansion.

EFT/factorization definition of renormalon: Asymptotic behavior of the perturbative expansion such that the associated ambiguity in the summation of the perturbative series can be absorbed into a higher order operator.

Example:

$$M_B = m_{\text{OS}} + \bar{\Lambda}_B + \mathcal{O}(1/m_{\text{OS}})$$

M_B is renormalon free. Therefore m_{OS} suffers from renormalon ambiguities:

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots)$$

with $B_n \sim n!$. In other words

$$\delta_{np}^{(\text{pert.})} m_{\text{OS}} = \delta_{np}^{(\text{pert.})} m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \sim \Lambda_{\text{QCD}}!$$

Modern view: EFT/Factorization

$$\mathcal{L} = \sum_n \frac{1}{m^n} c_n O_n \quad c(\nu) = \bar{c} + \sum_{n=0}^{\infty} c_n \alpha_s^{n+1}.$$

The Wilson coefficients are believed to be asymptotic: $c_n \sim n!$

IF SO such behavior should comply with the Operator Product Expansion.

EFT/factorization definition of renormalon: Asymptotic behavior of the perturbative expansion such that the associated ambiguity in the summation of the perturbative series can be absorbed into a higher order operator.

Example:

$$M_B = m_{\text{OS}} + \bar{\Lambda}_B + \mathcal{O}(1/m_{\text{OS}})$$

M_B is renormalon free. Therefore m_{OS} suffers from renormalon ambiguities:

$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots)$$

with $B_n \sim n!$. In other words

$$\delta_{np}^{(\text{pert.})} m_{\text{OS}} = \delta_{np}^{(\text{pert.})} m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \sim \Lambda_{\text{QCD}}!$$

Modern view: EFT/Factorization

$$\mathcal{L} = \sum_n \frac{1}{m^n} c_n O_n \quad c(\nu) = \bar{c} + \sum_{n=0}^{\infty} c_n \alpha_s^{n+1}.$$

The Wilson coefficients are believed to be asymptotic: $c_n \sim n!$

IF SO such behavior should comply with the Operator Product Expansion.

EFT/factorization definition of renormalon: Asymptotic behavior of the perturbative expansion such that the associated ambiguity in the summation of the perturbative series can be absorbed into a higher order operator.

Example:


$$M_B = m_{\text{OS}} + \bar{\Lambda}_B + \mathcal{O}(1/m_{\text{OS}})$$


M_B is renormalon free. Therefore m_{OS} suffers from renormalon ambiguities:


$$m_{\text{OS}} = m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots)$$

with $B_n \sim n!$. In other words

$$\delta_{np}^{(\text{pert.})} m_{\text{OS}} = \delta_{np}^{(\text{pert.})} m_{\overline{\text{MS}}} (1 + B_1 \alpha_s + B_2 \alpha_s^2 + \dots) \sim \Lambda_{\text{QCD}}!$$







Maximal accuracy of the Wilson coefficients from a perturbative calculation is (roughly) of the order of

$$\delta c \sim c_{n^*} \alpha_s^{n^*},$$

where $n^* \sim \frac{a}{\alpha_s}$. If a is positive c suffers from a non-perturbative ambiguity of order

$$\delta c \sim (\Lambda_{\text{QCD}})^{\frac{|a|\beta_0}{2\pi}}.$$

The Borel transform of $c(\nu)$ reads

$$B[c](t) \equiv \sum_{n=0}^{\infty} c_n \frac{t^n}{n!},$$

and c is written in terms of its Borel transform as

$$c = \bar{c} + \int_0^{\infty} dt e^{-t/\alpha_s} B[c](t).$$

The ambiguities in the Wilson coefficient ($c_n \sim n!$) reflects in poles in the Borel transform. If we take the one closest to the origin,

$$\delta B[c](t) \sim \frac{1}{a-t},$$

where a is a positive number.

Maximal accuracy of the Wilson coefficients from a perturbative calculation is (roughly) of the order of

$$\delta c \sim c_{n^*} \alpha_s^{n^*},$$

where $n^* \sim \frac{a}{\alpha_s}$. If a is positive c suffers from a non-perturbative ambiguity of order

$$\delta c \sim (\Lambda_{\text{QCD}})^{\frac{|a|\beta_0}{2\pi}}.$$

The Borel transform of $c(\nu)$ reads

$$B[c](t) \equiv \sum_{n=0}^{\infty} c_n \frac{t^n}{n!},$$

and c is written in terms of its Borel transform as

$$c = \bar{c} + \int_0^{\infty} dt e^{-t/\alpha_s} B[c](t).$$

The ambiguities in the Wilson coefficient ($c_n \sim n!$) reflects in poles in the Borel transform. If we take the one closest to the origin,

$$\delta B[c](t) \sim \frac{1}{a-t},$$

where a is a positive number.

Where do we expect renormalon effects to be more important?

Current-current correlator ($c_{n^*} \alpha_s^{n*+1} \sim \Lambda_{\text{QCD}}^4 / Q^4 \rightarrow c_n \sim n!$):

$$\int d^4x e^{iqx} \langle \text{vac} | J(x) J(0) | \text{vac} \rangle = (\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^4}{Q^4} + \dots$$

Plaquette:

$$\langle P \rangle = (\text{Pert. th.}) + a^4 \Lambda_{\text{QCD}}^4 + \dots$$

DIS:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^2}{Q^2} + \dots$$

Heavy quark physics:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}}{m_Q} + \dots$$

The natural place to look for these effects.

Where do we expect renormalon effects to be more important?

Current-current correlator ($c_{n^*} \alpha_s^{n^*+1} \sim \Lambda_{\text{QCD}}^4 / Q^4 \rightarrow c_n \sim n!$):

$$\int d^4x e^{iqx} \langle \text{vac} | J(x) J(0) | \text{vac} \rangle = (\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^4}{Q^4} + \dots$$

Plaquette:

$$\langle P \rangle = (\text{Pert. th.}) + a^4 \Lambda_{\text{QCD}}^4 + \dots$$

DIS:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^2}{Q^2} + \dots$$

Heavy quark physics:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}}{m_Q} + \dots$$

The natural place to look for these effects.

Where do we expect renormalon effects to be more important?

Current-current correlator ($c_{n^*} \alpha_s^{n*+1} \sim \Lambda_{\text{QCD}}^4 / Q^4 \rightarrow c_n \sim n!$):

$$\int d^4x e^{iqx} \langle \text{vac} | J(x) J(0) | \text{vac} \rangle = (\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^4}{Q^4} + \dots$$

Plaquette:

$$\langle P \rangle = (\text{Pert. th.}) + a^4 \Lambda_{\text{QCD}}^4 + \dots$$

DIS:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^2}{Q^2} + \dots$$

Heavy quark physics:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}}{m_Q} + \dots$$

The natural place to look for these effects.

Where do we expect renormalon effects to be more important?

Current-current correlator ($c_{n^*} \alpha_s^{n^*+1} \sim \Lambda_{\text{QCD}}^4 / Q^4 \rightarrow c_n \sim n!$):

$$\int d^4x e^{iqx} \langle \text{vac} | J(x) J(0) | \text{vac} \rangle = (\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^4}{Q^4} + \dots$$

Plaquette:

$$\langle P \rangle = (\text{Pert. th.}) + a^4 \Lambda_{\text{QCD}}^4 + \dots$$

DIS:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}^2}{Q^2} + \dots$$

Heavy quark physics:

$$(\text{Pert. th.}) + \frac{\Lambda_{\text{QCD}}}{m_Q} + \dots$$

The natural place to look for these effects.

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^{\infty} dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$(\text{Beneke}) \quad \text{Im} \int_0^{\infty} dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$

$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$\text{(Beneke)} \quad \text{Im} \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$

$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$\text{(Beneke)} \quad \text{Im} \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$

$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$(\text{Beneke}) \quad \text{Im} \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$

$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$(\text{Beneke}) \quad \text{Im} \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$

$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Pole mass: $m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1}$

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t), \quad B[m_{\text{OS}}](t) \equiv \sum_{n=0}^{\infty} r_n \frac{t^n}{n!}.$$

The large n dependence of r_n is dictated by the closest singularity to the origin of $B[m_{\text{OS}}]$ ($u = \frac{\beta_0 t}{4\pi}$).

$$(\text{Beneke}) \quad \text{Im} \int_0^\infty dt e^{-t/\alpha_s} B[m_{\text{OS}}](t) \sim \Lambda_{QCD} \sim \nu e^{-\frac{2\pi}{\beta_0 \alpha(\nu)}}$$





$$B[m_{\text{OS}}](t) = N_m \nu \frac{1}{(1-2u)^{1+b}} \left(1 + c_1(1-2u) + c_2(1-2u)^2 + \dots \right) + (\text{analytic term}),$$

$$r_n^{\text{asym}} = N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Beneke; Pineda

Check : $r_n \xrightarrow{n \rightarrow \infty} m_{\overline{\text{MS}}} \left(\frac{\beta_0}{2\pi} \right)^n n! N_m \sum_{s=0}^n \frac{\ln^s [\nu / m_{\overline{\text{MS}}}] }{s!} \sim \nu$



$\overline{\text{MS}}$: $\mathcal{O}(\alpha^4)$:

AP: hep-ph/0105008, hep-ph/0208031, hep-lat/0509022

Bali, AP: hep-ph/0310130

Ayala, Cvetic, AP, 1407.2128

Beneke, Marquard, Nason, Steinhauser, 1605.03609

....

We would like to have a proof (at the same level of existing proofs of a linear potential at long distances), beyond any reasonable doubt, of the existence of the renormalon in QCD.

Lattice: $\mathcal{O}(\alpha^{20})$:

Bali, Bauer, AP: arXiv:1111.3946

Bali, Bauer, AP, Torrero: arXiv:1303.3279

Bali, Bauer, AP: arXiv:1311.0114

$\overline{\text{MS}}$: $\mathcal{O}(\alpha^4)$:

AP: hep-ph/0105008, hep-ph/0208031, hep-lat/0509022

Bali, AP: hep-ph/0310130

Ayala, Cvetic, AP, 1407.2128

Beneke, Marquard, Nason, Steinhauser, 1605.03609

....

We would like to have a proof (at the same level of existing proofs of a linear potential at long distances), beyond any reasonable doubt, of the existence of the renormalon in QCD.

Lattice: $\mathcal{O}(\alpha^{20})$:

Bali, Bauer, AP: arXiv:1111.3946

Bali, Bauer, AP, Torrero: arXiv:1303.3279

Bali, Bauer, AP: arXiv:1311.0114

POLYAKOV LOOP versus δm (and m)

Possible to compute the energy of an static source in the lattice: δm of HQET.
We use Numerical Stochastic Perturbation Theory (Di Renzo et al.).

$$L^{(R)}(N_S, N_T) = \frac{1}{N_S^3} \sum_n \frac{1}{d_R} \text{tr} \left[\prod_{n_4=0}^{N_T-1} U_4^R(n) \right] \quad U_\mu^R(n) \approx e^{i A_\mu^R [(n+1/2)a]}$$

We implement triplet and octet representations R ($d_R = 3, 8$).

$$\langle L^{(R)}(N_S, N_T) \rangle \stackrel{N_T \rightarrow \infty}{\sim} e^{-N_T a \delta m^{(R)}(N_S)}$$

$$\delta m^{(R)}(N_S) = \lim_{N_T \rightarrow \infty} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S, N_T) \alpha^{n+1} = \frac{1}{a} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S) \alpha^{n+1} (1/a)$$

POLYAKOV LOOP versus δm (and m)

Possible to compute the energy of an static source in the lattice: δm of HQET.
We use Numerical Stochastic Perturbation Theory (Di Renzo et al.).

$$L^{(R)}(N_S, N_T) = \frac{1}{N_S^3} \sum_n \frac{1}{d_R} \text{tr} \left[\prod_{n_4=0}^{N_T-1} U_4^R(n) \right] \quad U_\mu^R(n) \approx e^{i A_\mu^R [(n+1/2)a]}$$

We implement triplet and octet representations R ($d_R = 3, 8$).

$$\langle L^{(R)}(N_S, N_T) \rangle \stackrel{N_T \rightarrow \infty}{\sim} e^{-N_T a \delta m^{(R)}(N_S)}$$

$$\delta m^{(R)}(N_S) = \lim_{N_T \rightarrow \infty} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S, N_T) \alpha^{n+1} = \frac{1}{a} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S) \alpha^{n+1} (1/a)$$

POLYAKOV LOOP versus δm (and m)

Possible to compute the energy of an static source in the lattice: δm of HQET.
We use Numerical Stochastic Perturbation Theory (Di Renzo et al.).

$$L^{(R)}(N_S, N_T) = \frac{1}{N_S^3} \sum_n \frac{1}{d_R} \text{tr} \left[\prod_{n_4=0}^{N_T-1} U_4^R(n) \right] \quad U_\mu^R(n) \approx e^{i A_\mu^R [(n+1/2)a]}$$

We implement triplet and octet representations R ($d_R = 3, 8$).

$$\langle L^{(R)}(N_S, N_T) \rangle \stackrel{N_T \rightarrow \infty}{\sim} e^{-N_T a \delta m^{(R)}(N_S)}$$

$$\delta m^{(R)}(N_S) = \lim_{N_T \rightarrow \infty} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S, N_T) \alpha^{n+1} = \frac{1}{a} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S) \alpha^{n+1} (1/a)$$

POLYAKOV LOOP versus δm (and m)

Possible to compute the energy of an static source in the lattice: δm of HQET.
We use Numerical Stochastic Perturbation Theory (Di Renzo et al.).

$$L^{(R)}(N_S, N_T) = \frac{1}{N_S^3} \sum_n \frac{1}{d_R} \text{tr} \left[\prod_{n_4=0}^{N_T-1} U_4^R(n) \right] \quad U_\mu^R(n) \approx e^{i A_\mu^R [(n+1/2)a]}$$

We implement triplet and octet representations R ($d_R = 3, 8$).

$$\langle L^{(R)}(N_S, N_T) \rangle \stackrel{N_T \rightarrow \infty}{\sim} e^{-N_T a \delta m^{(R)}(N_S)}$$

$$\delta m^{(R)}(N_S) = \lim_{N_T \rightarrow \infty} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S, N_T) \alpha^{n+1} = \frac{1}{a} \sum_{n=0}^{\infty} c_n^{(R,\rho)}(N_S) \alpha^{n+1} (1/a)$$

Perturbative OPE (Zimmermann) at finite volume ($N_S \rightarrow \infty$)

$$\delta m = \lim_{N_S \rightarrow \infty} \delta m(N_S) \quad c_n = \lim_{N_S \rightarrow \infty} c_n(N_S) \quad \left(\lim_{n \rightarrow \infty} c_n^{(3,\rho)} = r_n(\nu)/\nu \right).$$

For large N_S , we write (OPE: $\frac{1}{a} \gg \frac{1}{N_S a}$)

$$\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} \left(a^{-1} \right) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} \left((aN_S)^{-1} \right) + \mathcal{O} \left(\frac{1}{N_S^2} \right).$$

Taylor expansion of $\alpha((aN_S)^{-1})$ in powers of $\alpha(a^{-1})$:

$$c_n(N_S) = c_n - \frac{f_n(N_S)}{N_S} + \mathcal{O} \left(\frac{1}{N_S^2} \right); \quad f_n(N_S) = \sum_{i=0}^n f_n^{(i)} \ln^i(N_S),$$

$f_n^{(0)} = f_n$. The coefficients $f_n^{(i)}$ for $i > 0$ are determined by f_m and β_j .

$$f_1(N_S) = f_1 + f_0 \frac{\beta_0}{2\pi} \ln(N_S),$$

$$f_2(N_S) = f_2 + \left[2f_1 \frac{\beta_0}{2\pi} + f_0 \frac{\beta_1}{8\pi^2} \right] \ln(N_S) + f_0 \left(\frac{\beta_0}{2\pi} \right)^2 \ln^2(N_S),$$

and so on.

Perturbative OPE (Zimmermann) at finite volume ($N_S \rightarrow \infty$)

$$\delta m = \lim_{N_S \rightarrow \infty} \delta m(N_S) \quad c_n = \lim_{N_S \rightarrow \infty} c_n(N_S) \quad \left(\lim_{n \rightarrow \infty} c_n^{(3,\rho)} = r_n(\nu)/\nu \right).$$

For large N_S , we write (OPE: $\frac{1}{a} \gg \frac{1}{N_S a}$)

$$\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} \left(a^{-1} \right) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} \left((aN_S)^{-1} \right) + \mathcal{O} \left(\frac{1}{N_S^2} \right).$$

Taylor expansion of $\alpha((aN_S)^{-1})$ in powers of $\alpha(a^{-1})$:

$$c_n(N_S) = c_n - \frac{f_n(N_S)}{N_S} + \mathcal{O} \left(\frac{1}{N_S^2} \right); \quad f_n(N_S) = \sum_{i=0}^n f_n^{(i)} \ln^i(N_S),$$

$f_n^{(0)} = f_n$. The coefficients $f_n^{(i)}$ for $i > 0$ are determined by f_m and β_j .

$$f_1(N_S) = f_1 + f_0 \frac{\beta_0}{2\pi} \ln(N_S),$$

$$f_2(N_S) = f_2 + \left[2f_1 \frac{\beta_0}{2\pi} + f_0 \frac{\beta_1}{8\pi^2} \right] \ln(N_S) + f_0 \left(\frac{\beta_0}{2\pi} \right)^2 \ln^2(N_S),$$

and so on.

Perturbative OPE (Zimmermann) at finite volume ($N_S \rightarrow \infty$)

$$\delta m = \lim_{N_S \rightarrow \infty} \delta m(N_S) \quad c_n = \lim_{N_S \rightarrow \infty} c_n(N_S) \quad \left(\lim_{n \rightarrow \infty} c_n^{(3,\rho)} = r_n(\nu)/\nu \right).$$

For large N_S , we write (OPE: $\frac{1}{a} \gg \frac{1}{N_S a}$)

$$\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} \left(a^{-1} \right) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} \left((aN_S)^{-1} \right) + \mathcal{O} \left(\frac{1}{N_S^2} \right).$$

Taylor expansion of $\alpha((aN_S)^{-1})$ in powers of $\alpha(a^{-1})$:

$$c_n(N_S) = c_n - \frac{f_n(N_S)}{N_S} + \mathcal{O} \left(\frac{1}{N_S^2} \right); \quad f_n(N_S) = \sum_{i=0}^n f_n^{(i)} \ln^i(N_S),$$

$f_n^{(0)} = f_n$. The coefficients $f_n^{(i)}$ for $i > 0$ are determined by f_m and β_j .

$$f_1(N_S) = f_1 + f_0 \frac{\beta_0}{2\pi} \ln(N_S),$$

$$f_2(N_S) = f_2 + \left[2f_1 \frac{\beta_0}{2\pi} + f_0 \frac{\beta_1}{8\pi^2} \right] \ln(N_S) + f_0 \left(\frac{\beta_0}{2\pi} \right)^2 \ln^2(N_S),$$

and so on.

"Physical interpretation"





Figure : *Self-interactions with replicas producing $1/L = 1/(aN_S)$ Coulomb terms.*


$$\delta m^{(R)}(N_S) \propto \int_{1/(aN_S)}^{1/a} dk \alpha(k) \sim \frac{1}{a} \sum_n c_n \alpha^{n+1}(a^{-1}) - \frac{1}{aN_S} \sum_n c_n \alpha^{n+1}((aN_S)^{-1}),$$


$$c_n \simeq N_m \left(\frac{\beta_0}{2\pi}\right)^n n!, \quad f_n^{(i)}(N_S) \simeq N_m \left(\frac{\beta_0}{2\pi}\right)^n \frac{n!}{i!}.$$

	$c_n^{(3,0)}$	$c_n^{(3,1/6)}$	$c_n^{(8,0)} C_F / C_A$	$c_n^{(8,1/6)} C_F / C_A$
c_0	2.117274357	0.72181(99)	2.117274357	0.72181(99)
c_1	11.136(11)	6.385(10)	11.140(12)	6.387(10)
$c_2/10$	8.610(13)	8.124(12)	8.587(14)	8.129(12)
$c_3/10^2$	7.945(16)	7.670(13)	7.917(20)	7.682(15)
$c_4/10^3$	8.215(34)	8.017(33)	8.197(42)	8.017(36)
$c_5/10^4$	9.322(59)	9.160(59)	9.295(76)	9.139(64)
$c_6/10^6$	1.153(11)	1.138(11)	1.144(13)	1.134(12)
$c_7/10^7$	1.558(21)	1.541(22)	1.533(25)	1.535(22)
$c_8/10^8$	2.304(43)	2.284(45)	2.254(51)	2.275(45)
$c_9/10^9$	3.747(95)	3.717(97)	3.64(11)	3.703(98)
$c_{10}/10^{10}$	6.70(22)	6.65(22)	6.49(25)	6.63(22)
$c_{11}/10^{12}$	1.316(52)	1.306(53)	1.269(59)	1.303(53)
$c_{12}/10^{13}$	2.81(13)	2.79(13)	2.71(14)	2.78(13)
$c_{13}/10^{14}$	6.51(35)	6.46(35)	6.29(37)	6.45(35)
$c_{14}/10^{16}$	1.628(96)	1.613(97)	1.57(10)	1.614(97)
$c_{15}/10^{17}$	4.36(28)	4.32(28)	4.22(29)	4.33(28)
$c_{16}/10^{19}$	1.247(86)	1.235(86)	1.206(89)	1.236(86)
$c_{17}/10^{20}$	3.78(28)	3.75(28)	3.66(28)	3.75(28)
$c_{18}/10^{22}$	1.215(93)	1.204(94)	1.176(95)	1.205(94)
$c_{19}/10^{23}$	4.12(33)	4.08(33)	3.99(34)	4.08(33)

	$f_n^{(3,0)}$	$f_n^{(3,1/6)}$	$f_n^{(8,0)} C_F / C_A$	$f_n^{(8,1/6)} C_F / C_A$
f_0	0.7696256328	0.7810(59)	0.7696256328	0.7810(69)
f_1	6.075(78)	6.046(58)	6.124(87)	6.063(68)
$f_2/10$	5.628(91)	5.644(62)	5.60(11)	5.691(78)
$f_3/10^2$	5.87(11)	5.858(76)	6.00(18)	5.946(91)
$f_4/10^3$	6.33(22)	6.29(17)	6.57(40)	6.26(23)
$f_5/10^4$	7.73(35)	7.71(26)	7.67(66)	7.78(42)
$f_6/10^5$	9.86(53)	9.80(42)	9.68(99)	9.79(69)
$f_7/10^7$	1.388(81)	1.378(71)	1.35(15)	1.38(11)
$f_8/10^8$	2.12(12)	2.11(12)	2.06(22)	2.10(17)
$f_9/10^9$	3.54(20)	3.52(20)	3.40(37)	3.51(27)
$f_{10}/10^{10}$	6.49(33)	6.44(34)	6.23(67)	6.44(43)
$f_{11}/10^{12}$	1.296(64)	1.286(66)	1.24(13)	1.286(74)
$f_{12}/10^{13}$	2.68(19)	2.64(18)	2.65(33)	2.65(21)
$f_{13}/10^{14}$	6.70(54)	6.68(52)	6.36(90)	6.66(57)
$f_{14}/10^{16}$	1.58(14)	1.56(14)	1.55(22)	1.57(15)
$f_{15}/10^{17}$	4.41(34)	4.37(33)	4.24(47)	4.37(35)
$f_{16}/10^{19}$	1.241(92)	1.230(91)	1.20(11)	1.231(94)
$f_{17}/10^{20}$	3.79(28)	3.75(28)	3.67(30)	3.76(28)
$f_{18}/10^{22}$	1.215(94)	1.204(94)	1.176(97)	1.205(94)
$f_{19}/10^{23}$	4.12(33)	4.08(33)	3.99(34)	4.08(33)










Figure : c_n times $\sqrt{n_0}$, for five different values of the lattice scheme coupling constant α , ranging from $\alpha(\nu) \approx 0.096$ ($n_0 = 5$) to $\alpha(\nu) \approx 0.036$ ($n_0 = 15$). Bali, Bauer, AP, Torrero, 1303.3279.

Ratios

$$\begin{aligned} \frac{c_n^{(3,\rho)} 1}{c_{n-1}^{(3,\rho)} n} &= \frac{c_n^{(8,\rho)} 1}{c_{n-1}^{(8,\rho)} n} \\ &= \frac{\beta_0}{2\pi} \left\{ 1 + \frac{b}{n} - \frac{bs_1}{n^2} + \frac{1}{n^3} [b^2 s_1^2 + b(b-1)(s_1 - 2s_2)] + \mathcal{O}\left(\frac{1}{n^4}\right) \right\} . \end{aligned}$$




Figure : Ratios $c_n/(n c_{n-1})$ of the smeared (blue) and unsmeared (red) triplet static self-energy coefficients c_n in comparison to the theoretical prediction at different orders in the $1/n$ expansion.




Figure : The ratios $c_n/(nc_{n-1})$ for the smeared and unsmeared, triplet and octet static self-energies, compared to the prediction for the LO, next-to-leading order (NLO), NNLO and NNNLO of the $1/n$ expansion.

N_m

$$c_n^{fitted} = N_m \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$f_n^{fitted} = N_m \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$




Figure : N_m^{latt} determined via r_n truncated at NNNLO, from the coefficients $c_n^{(3,0)}$, $c_n^{(3,1/6)}$, $f_n^{(3,0)}$ and $f_n^{(3,1/6)}$. The horizontal band is our final result: $N_m = 17.9(1.0)$

From lattice to $\overline{\text{MS}}$ scheme

$$\alpha_{\overline{\text{MS}}}(\mu) = \alpha_{\text{latt}}(\mu) \left(1 + d_1 \alpha_{\text{latt}}(\mu) + d_2 \alpha_{\text{latt}}^2(\mu) + d_3 \alpha_{\text{latt}}^3(\mu) + \mathcal{O}(\alpha_{\text{latt}}^4) \right),$$

$$N_{m, m_{\tilde{g}}}^{\overline{\text{MS}}} = N_{m, m_{\tilde{g}}}^{\text{latt}} \Lambda_{\text{latt}} / \Lambda_{\overline{\text{MS}}}, \quad \text{where} \quad \Lambda_{\overline{\text{MS}}} = e^{\frac{2\pi d_1}{\beta_0}} \Lambda_{\text{latt}} \approx 28.809338139488 \Lambda_{\text{latt}}.$$

This yields the numerical values

$$N_m^{\overline{\text{MS}}} = 0.620(35), \quad C_F/C_A N_{m_{\tilde{g}}}^{\overline{\text{MS}}} = -C_F/C_A N_{\Lambda}^{\overline{\text{MS}}} = 0.610(41).$$

~ 20 standard deviations from zero!

From $N_m^{\overline{\text{MS}}} = 0.600(29)$ (Ayala, Cvetic, AP). Combined $N_m^{\overline{\text{MS}}} = 0.608(22)$.

From lattice to $\overline{\text{MS}}$ scheme

$$\alpha_{\overline{\text{MS}}}(\mu) = \alpha_{\text{latt}}(\mu) \left(1 + d_1 \alpha_{\text{latt}}(\mu) + d_2 \alpha_{\text{latt}}^2(\mu) + d_3 \alpha_{\text{latt}}^3(\mu) + \mathcal{O}(\alpha_{\text{latt}}^4) \right),$$

$$N_{m, m_{\tilde{g}}}^{\overline{\text{MS}}} = N_{m, m_{\tilde{g}}}^{\text{latt}} \Lambda_{\text{latt}} / \Lambda_{\overline{\text{MS}}}, \quad \text{where} \quad \Lambda_{\overline{\text{MS}}} = e^{\frac{2\pi d_1}{\beta_0}} \Lambda_{\text{latt}} \approx 28.809338139488 \Lambda_{\text{latt}}.$$

This yields the numerical values

$$N_m^{\overline{\text{MS}}} = 0.620(35), \quad C_F/C_A N_{m_{\tilde{g}}}^{\overline{\text{MS}}} = -C_F/C_A N_{\Lambda}^{\overline{\text{MS}}} = 0.610(41).$$

~ 20 standard deviations from zero!

From $N_m^{\overline{\text{MS}}} = 0.600(29)$ (Ayala, Cvetic, AP). Combined $N_m^{\overline{\text{MS}}} = 0.608(22)$.

From lattice to $\overline{\text{MS}}$ scheme

$$\alpha_{\overline{\text{MS}}}(\mu) = \alpha_{\text{latt}}(\mu) \left(1 + d_1 \alpha_{\text{latt}}(\mu) + d_2 \alpha_{\text{latt}}^2(\mu) + d_3 \alpha_{\text{latt}}^3(\mu) + \mathcal{O}(\alpha_{\text{latt}}^4) \right),$$

$$N_{m, m_{\tilde{g}}}^{\overline{\text{MS}}} = N_{m, m_{\tilde{g}}}^{\text{latt}} \Lambda_{\text{latt}} / \Lambda_{\overline{\text{MS}}}, \quad \text{where} \quad \Lambda_{\overline{\text{MS}}} = e^{\frac{2\pi d_1}{\beta_0}} \Lambda_{\text{latt}} \approx 28.809338139488 \Lambda_{\text{latt}}.$$

This yields the numerical values

$$N_m^{\overline{\text{MS}}} = 0.620(35), \quad C_F/C_A N_{m_{\tilde{g}}}^{\overline{\text{MS}}} = -C_F/C_A N_{\Lambda}^{\overline{\text{MS}}} = 0.610(41).$$

~ 20 standard deviations from zero!

From $N_m^{\overline{\text{MS}}} = 0.600(29)$ (Ayala, Cvetic, AP). Combined $N_m^{\overline{\text{MS}}} = 0.608(22)$.

Plaquette (Bali, Bauer, AP: 1401.7999, 1403.6477)

$$\langle P \rangle_{\text{pert}}(N) \equiv \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = \sum_{n \geq 0} p_n(N) \alpha^{n+1}$$

Perturbative OPE

$$\frac{1}{a} \gg \frac{1}{Na} \rightarrow \langle P \rangle_{\text{pert}}(N) = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{soft}} + \mathcal{O}\left(\frac{1}{N^6}\right),$$

where

$$P_{\text{pert}} = \sum_{n \geq 0} p_n \alpha^{n+1}, \quad C_G = 1 + \sum_{k \geq 0} c_k \alpha^{k+1}, \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

$$\begin{aligned} \langle P \rangle_{\text{pert}}(N) &= \sum_{n \geq 0} \left[p_n - \frac{f_n(N)}{N^4} \right] \alpha^{n+1} \\ &= \sum_{n \geq 0} p_n \alpha^{n+1} - \frac{1}{N^4} \left(1 + \sum_{k \geq 0} c_k \alpha^{k+1} (a^{-1}) \right) \times \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1}) + \mathcal{O}\left(\frac{1}{N^6}\right), \end{aligned}$$

$$\left(\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} (a^{-1}) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} ((aN_S)^{-1}) + \mathcal{O}\left(\frac{1}{N_S^2}\right) \right)$$

Plaquette (Bali, Bauer, AP: 1401.7999, 1403.6477)

$$\langle P \rangle_{\text{pert}}(N) \equiv \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = \sum_{n \geq 0} p_n(N) \alpha^{n+1}$$

Perturbative OPE

$$\frac{1}{a} \gg \frac{1}{Na} \rightarrow \langle P \rangle_{\text{pert}}(N) = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{soft}} + \mathcal{O}\left(\frac{1}{N^6}\right),$$

where

$$P_{\text{pert}} = \sum_{n \geq 0} p_n \alpha^{n+1}, \quad C_G = 1 + \sum_{k \geq 0} c_k \alpha^{k+1}, \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

$$\begin{aligned} \langle P \rangle_{\text{pert}}(N) &= \sum_{n \geq 0} \left[p_n - \frac{f_n(N)}{N^4} \right] \alpha^{n+1} \\ &= \sum_{n \geq 0} p_n \alpha^{n+1} - \frac{1}{N^4} \left(1 + \sum_{k \geq 0} c_k \alpha^{k+1} (a^{-1}) \right) \times \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1}) + \mathcal{O}\left(\frac{1}{N^6}\right), \end{aligned}$$

$$\left(\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} (a^{-1}) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} ((aN_S)^{-1}) + \mathcal{O}\left(\frac{1}{N_S^2}\right) \right)$$

Plaquette (Bali, Bauer, AP: 1401.7999, 1403.6477)

$$\langle P \rangle_{\text{pert}}(N) \equiv \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = \sum_{n \geq 0} p_n(N) \alpha^{n+1}$$

Perturbative OPE

$$\frac{1}{a} \gg \frac{1}{Na} \rightarrow \langle P \rangle_{\text{pert}}(N) = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{soft}} + \mathcal{O}\left(\frac{1}{N^6}\right),$$

where

$$P_{\text{pert}} = \sum_{n \geq 0} p_n \alpha^{n+1}, \quad C_G = 1 + \sum_{k \geq 0} c_k \alpha^{k+1}, \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

$$\begin{aligned} \langle P \rangle_{\text{pert}}(N) &= \sum_{n \geq 0} \left[p_n - \frac{f_n(N)}{N^4} \right] \alpha^{n+1} \\ &= \sum_{n \geq 0} p_n \alpha^{n+1} - \frac{1}{N^4} \left(1 + \sum_{k \geq 0} c_k \alpha^{k+1} (a^{-1}) \right) \times \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1}) + \mathcal{O}\left(\frac{1}{N^6}\right), \\ \left(\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} (a^{-1}) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} ((aN_S)^{-1}) + \mathcal{O}\left(\frac{1}{N_S^2}\right) \right) \end{aligned}$$

Plaquette (Bali, Bauer, AP: 1401.7999, 1403.6477)

$$\langle P \rangle_{\text{pert}}(N) \equiv \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = \sum_{n \geq 0} p_n(N) \alpha^{n+1}$$

Perturbative OPE

$$\frac{1}{a} \gg \frac{1}{Na} \rightarrow \langle P \rangle_{\text{pert}}(N) = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{soft}} + \mathcal{O}\left(\frac{1}{N^6}\right),$$

where

$$P_{\text{pert}} = \sum_{n \geq 0} p_n \alpha^{n+1}, \quad C_G = 1 + \sum_{k \geq 0} c_k \alpha^{k+1}, \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

$$\begin{aligned} \langle P \rangle_{\text{pert}}(N) &= \sum_{n \geq 0} \left[p_n - \frac{f_n(N)}{N^4} \right] \alpha^{n+1} \\ &= \sum_{n \geq 0} p_n \alpha^{n+1} - \frac{1}{N^4} \left(1 + \sum_{k \geq 0} c_k \alpha^{k+1} (a^{-1}) \right) \times \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1}) + \mathcal{O}\left(\frac{1}{N^6}\right), \\ \left(\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} (a^{-1}) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} ((aN_S)^{-1}) + \mathcal{O}\left(\frac{1}{N_S^2}\right) \right) \end{aligned}$$

Plaquette (Bali, Bauer, AP: 1401.7999, 1403.6477)

$$\langle P \rangle_{\text{pert}}(N) \equiv \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = \sum_{n \geq 0} p_n(N) \alpha^{n+1}$$

Perturbative OPE

$$\frac{1}{a} \gg \frac{1}{Na} \rightarrow \langle P \rangle_{\text{pert}}(N) = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{soft}} + \mathcal{O}\left(\frac{1}{N^6}\right),$$

where

$$P_{\text{pert}} = \sum_{n \geq 0} p_n \alpha^{n+1}, \quad C_G = 1 + \sum_{k \geq 0} c_k \alpha^{k+1}, \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

$$\begin{aligned} \langle P \rangle_{\text{pert}}(N) &= \sum_{n \geq 0} \left[p_n - \frac{f_n(N)}{N^4} \right] \alpha^{n+1} \\ &= \sum_{n \geq 0} p_n \alpha^{n+1} - \frac{1}{N^4} \left(1 + \sum_{k \geq 0} c_k \alpha^{k+1} (a^{-1}) \right) \times \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1}) + \mathcal{O}\left(\frac{1}{N^6}\right), \\ \left(\delta m(N_S) = \frac{1}{a} \sum_{n=0}^{\infty} c_n \alpha^{n+1} (a^{-1}) - \frac{1}{aN_S} \sum_{n=0}^{\infty} f_n \alpha^{n+1} ((aN_S)^{-1}) + \mathcal{O}\left(\frac{1}{N_S^2}\right) \right) \end{aligned}$$

$$\langle P \rangle = \sum_{n=0}^N p_n \alpha^{n+1}(a^{-1}) + a^4 \frac{\pi^2}{36} \langle G^2 \rangle + \dots$$

$$d = 1 (n_0 \sim 7) \longrightarrow d = 4 (n_0 \sim 28)$$

$$N+1 = 35$$

(before Di Renzo et al. N+1=8; Horsley et al. N+1=20)
 Renormalon expectations:

$$p_n^{\text{latt}} \stackrel{n \rightarrow \infty}{=} N_P^{\text{latt}} \left(\frac{\beta_0}{2\pi d} \right)^n \frac{\Gamma(n+1+db)}{\Gamma(1+db)} \left\{ 1 + \frac{20.09}{n+db} + \frac{505 \pm 33}{(n+db)^2} + \mathcal{O}\left(\frac{1}{n^3}\right) \right\}.$$

$$\frac{p_n}{np_{n-1}} = \frac{\beta_0}{2\pi d} \left\{ 1 + \frac{db}{n} + \frac{db(1 - ds_1)}{n^2} + \frac{\#}{n^3} + \mathcal{O}\left(\frac{1}{n^4}\right) \right\}.$$

$$\langle P \rangle = \sum_{n=0}^N p_n \alpha^{n+1} (a^{-1}) + a^4 \frac{\pi^2}{36} \langle G^2 \rangle + \dots$$

$$d = 1 (n_0 \sim 7) \longrightarrow d = 4 (n_0 \sim 28)$$

$$N+1 = 35$$

(before Di Renzo et al. N+1=8; Horsley et al. N+1=20)

Renormalon expectations:

$$p_n^{\text{latt}} \stackrel{n \rightarrow \infty}{=} N_P^{\text{latt}} \left(\frac{\beta_0}{2\pi d} \right)^n \frac{\Gamma(n+1+db)}{\Gamma(1+db)} \left\{ 1 + \frac{20.09}{n+db} + \frac{505 \pm 33}{(n+db)^2} + \mathcal{O}\left(\frac{1}{n^3}\right) \right\}.$$

$$\frac{p_n}{np_{n-1}} = \frac{\beta_0}{2\pi d} \left\{ 1 + \frac{db}{n} + \frac{db(1 - ds_1)}{n^2} + \frac{\#}{n^3} + \mathcal{O}\left(\frac{1}{n^4}\right) \right\}.$$

$$\langle P \rangle = \sum_{n=0}^N p_n \alpha^{n+1} (a^{-1}) + a^4 \frac{\pi^2}{36} \langle G^2 \rangle + \dots$$

$$d = 1 (n_0 \sim 7) \longrightarrow d = 4 (n_0 \sim 28)$$

$$N+1 = 35$$

(before Di Renzo et al. N+1=8; Horsley et al. N+1=20)

Renormalon expectations:

$$p_n^{\text{latt}} \stackrel{n \rightarrow \infty}{=} N_P^{\text{latt}} \left(\frac{\beta_0}{2\pi d} \right)^n \frac{\Gamma(n+1+db)}{\Gamma(1+db)} \left\{ 1 + \frac{20.09}{n+db} + \frac{505 \pm 33}{(n+db)^2} + \mathcal{O}\left(\frac{1}{n^3}\right) \right\} .$$

$$\frac{p_n}{np_{n-1}} = \frac{\beta_0}{2\pi d} \left\{ 1 + \frac{db}{n} + \frac{db(1 - ds_1)}{n^2} + \frac{\#}{n^3} + \mathcal{O}\left(\frac{1}{n^4}\right) \right\} .$$




Figure : Ratios $p_n / (np_{n-1})$ of the plaquette coefficients p_n ($N = \infty$, $N = 28$) in comparison to the theoretical prediction at different orders in the $1/n$ expansion.




Figure : N_P , determined from the coefficients p_n truncated at NLO, NNLO and NNNLO. The green box marks our final result.

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$




Figure : N_P , determined from the coefficients p_n truncated at NLO, NNLO and NNNLO. The green box marks our final result.

$$N_P^{\overline{MS}} = 0.61(25) \quad N_G^{\overline{MS}} = \frac{36}{\pi^2} N_P^{\overline{MS}} = 2.24(92).$$

Uncertainty of the sum due to the truncation

$$\delta S_P = \sqrt{n_0} p_{n_0} \alpha^{n_0+1} \approx \frac{(2\pi)^{3/2} d^{1+db}}{2^{db} \beta_0 \Gamma(1+db)} N_P(\Lambda a)^4 \approx 12.06 N_P(\Lambda a)^4.$$

This object is scheme- and scale-independent (to $1/n$ -precision)




Figure : $\sqrt{n} p_n \alpha^{n+1} / (\Lambda_{latt} a)^4$, versus n for $\beta = 5.3, 5.8, 6.3, 6.8$ and 7.3 . The green band is the theoretical expectation $12.06 N_P = 5.1(2.1) \times 10^6$.

$$\sqrt{n_0} \frac{|r_{n_0}|}{\Lambda_{\text{latt}}} \alpha^{n_0+1}(\nu) = \frac{2^{3/2-b} \pi^{3/2}}{\beta_0 \Gamma(1+b)} |N_m| \approx 1.206 |N_m|,$$




Figure : c_n times $\sqrt{n_0}$, for five different values of the lattice scheme coupling constant α , ranging from $\alpha(\nu) \approx 0.096$ ($n_0 = 5$) to $\alpha(\nu) \approx 0.036$ ($n_0 = 15$).

ONLY once the asymptotic behavior of perturbation theory has been reached one can go beyond perturbation theory (at last...)

BUT NOT TODAY

ONLY once the asymptotic behavior of perturbation theory has been reached one can go beyond perturbation theory (at last...)

BUT NOT TODAY

$$\delta \langle G^2 \rangle_{\text{NP}} \simeq \frac{(2\pi)^{3/2} d^{1+db}}{2^{db} \beta_0 \Gamma(1+db)} N_G^{\overline{\text{MS}}} \Big|_{n_f=0} \Lambda_{\overline{\text{MS}}}^4 = 27(11) \Lambda_{\overline{\text{MS}}}^4 \sim 0.087 \text{ GeV}^4.$$

$$\langle G^2 \rangle = 3.18(29) r_0^{-4} = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4})$ $\overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4}) \overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4})$ $\overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4}) \overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4}) \overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4})$ $\overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS

Renormalons go beyond large- β_0 analysis: \rightarrow (NP)OPE

Strong evidence of renormalon dominance in heavy quark physics from $\mathcal{O}(\alpha^{3/4})$ $\overline{\text{MS}}$ -like computations: Pole mass, static potential, ...

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.600(29), \quad N_m^{\overline{\text{MS}}}(n_l = 3) = 0.563(26).$$

Lattice: For the first time it was possible to follow the factorial growth of the coefficients over many orders, from around α^9 up to α^{20} , vastly increasing the credibility of the prediction.

$$N_m^{\overline{\text{MS}}}(n_l = 0) = 0.620(35), \quad C_F/C_A N_\Lambda^{\overline{\text{MS}}}(n_l = 0) = -0.610(41).$$

Two independent determinations with very different systematics.

We have (numerically) proven, beyond any reasonable doubt (~ 20 standard deviations!), the existence of the renormalon in QCD.

RENORMALON OK

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

CONCLUSIONS: OPE and the plaquette

$$N_P^{\overline{\text{MS}}} = 0.61(25) \quad N_G^{\overline{\text{MS}}} = \frac{36}{\pi^2} N_P^{\overline{\text{MS}}} = 2.24(92).$$

Nonperturbative quantities ($\bar{\Lambda}$, Λ_H , $\langle G^2 \rangle$, ...) can only be defined after subtracting the divergent perturbative series.

$$\delta \langle G^2 \rangle_{\text{NP}} = 27(11) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.087 \text{ GeV}^4. \quad \langle G^2 \rangle = 24.2(8.0) \Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$

Non-perturbative OPE OK (for the plaquette)

Dimension two condensates: artifacts of incomplete subtractions

- ▶ unquantifiable error due to the simplified parameterization of higher order perturbation theory
- ▶ short distance effect → process dependent

FUTURE:

n_f dependence

Control of the subtraction-scheme dependence?

Problem for sum rules?

Lattice → Model independent/systematic procedure to get ALL condensates

Beyond perturbation theory (at last...)

$$\langle P \rangle_{\text{pert}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle O_G \rangle_{\text{soft}} + \mathcal{O}(a^6).$$

$$\frac{1}{a} \gg \frac{1}{Na}$$

$$\langle P \rangle_{\text{MC}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{MC}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{MC}} + \mathcal{O}(a^6).$$

$$\frac{1}{a} \gg \frac{1}{Na} \gg \Lambda_{\text{QCD}} \rightarrow \langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{soft}} \left[1 + \mathcal{O}(\Lambda_{\text{QCD}}^2 (Na)^2) \right]$$

$$\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na} \rightarrow \langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{NP}} \left[1 + \mathcal{O} \left(\frac{1}{\Lambda_{\text{QCD}}^2 (Na)^2} \right) \right],$$

where $\langle G^2 \rangle_{\text{NP}} \sim \Lambda_{\text{QCD}}^4$ is the NP gluon condensate (Vainshtein, Zakharov, Shifman).

Beyond perturbation theory (at last...)

$$\langle P \rangle_{\text{pert}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{NSPT}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle O_G \rangle_{\text{soft}} + \mathcal{O}(a^6).$$

$$\frac{1}{a} \gg \frac{1}{Na}$$

$$\langle P \rangle_{\text{MC}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{MC}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{MC}} + \mathcal{O}(a^6).$$

$$\frac{1}{a} \gg \frac{1}{Na} \gg \Lambda_{\text{QCD}} \rightarrow \langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{soft}} \left[1 + \mathcal{O}(\Lambda_{\text{QCD}}^2 (Na)^2) \right]$$

$$\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na} \rightarrow \langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{NP}} \left[1 + \mathcal{O} \left(\frac{1}{\Lambda_{\text{QCD}}^2 (Na)^2} \right) \right],$$

where $\langle G^2 \rangle_{\text{NP}} \sim \Lambda_{\text{QCD}}^4$ is the NP gluon condensate (Vainshtein, Zakharov, Shifman).

$$\frac{1}{a} \gg \frac{1}{Na} \gg \Lambda_{\text{QCD}}$$

$$\langle P \rangle_{\text{MC}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{MC}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{MC}} + \mathcal{O}(a^6) .$$

$$\langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{soft}} \left[1 + \mathcal{O}(\Lambda_{\text{QCD}}^2 (Na)^2) \right], \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

In this limit non-perturbative effects can be computed at weak coupling (still far from trivial: see resurgence analysis in 1+1 dimensions).

Observations:

- ▶ In this limit the gluon condensate renormalon is not produced by non-perturbative effects.
- ▶ The resummation of all $(\Lambda_{\text{QCD}}^2 (Na)^2)^n$ effects remains to be done to reach the scaling region at infinite volume: $\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na}$, i.e. $\langle G^2 \rangle_{\text{NP}}$.

$$\frac{1}{a} \gg \frac{1}{Na} \gg \Lambda_{\text{QCD}}$$

$$\langle P \rangle_{\text{MC}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{MC}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{MC}} + \mathcal{O}(a^6) .$$

$$\langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{soft}} \left[1 + \mathcal{O}(\Lambda_{\text{QCD}}^2 (Na)^2) \right], \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

In this limit non-perturbative effects can be computed at weak coupling (still far from trivial: see resurgence analysis in 1+1 dimensions).

Observations:

- ▶ In this limit the gluon condensate renormalon is not produced by non-perturbative effects.
- ▶ The resummation of all $(\Lambda_{\text{QCD}}^2 (Na)^2)^n$ effects remains to be done to reach the scaling region at infinite volume: $\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na}$, i.e. $\langle G^2 \rangle_{\text{NP}}$.

$$\frac{1}{a} \gg \frac{1}{Na} \gg \Lambda_{\text{QCD}}$$

$$\langle P \rangle_{\text{MC}} = \frac{1}{Z} \int [dU_{x,\mu}] e^{-S[U]} P[U] \Big|_{\text{MC}} = P_{\text{pert}}(\alpha) \langle 1 \rangle + \frac{\pi^2}{36} C_G(\alpha) a^4 \langle G^2 \rangle_{\text{MC}} + \mathcal{O}(a^6) .$$

$$\langle G^2 \rangle_{\text{MC}} = \langle G^2 \rangle_{\text{soft}} \left[1 + \mathcal{O}(\Lambda_{\text{QCD}}^2 (Na)^2) \right], \quad \frac{\pi^2}{36} a^4 \langle G^2 \rangle_{\text{soft}} = -\frac{1}{N^4} \sum_{n \geq 0} f_n \alpha^{n+1} ((Na)^{-1})$$

In this limit non-perturbative effects can be computed at weak coupling (still far from trivial: see resurgence analysis in 1+1 dimensions).

Observations:

- ▶ In this limit the gluon condensate renormalon is not produced by non-perturbative effects.
- ▶ The resummation of all $(\Lambda_{\text{QCD}}^2 (Na)^2)^n$ effects remains to be done to reach the scaling region at infinite volume: $\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na}$, i.e. $\langle G^2 \rangle_{\text{NP}}$.

Determination of the gluon condensate: $\frac{1}{a} \gg \Lambda_{\text{QCD}} \gg \frac{1}{Na}$

$$\langle G^2 \rangle_{\text{NP}} = \frac{36 C_G^{-1}(\alpha)}{\pi^2 a^4(\alpha)} [\langle P \rangle_{\text{MC}}(\alpha) - S_P(\alpha)] + \mathcal{O}(a^2 \Lambda_{\text{QCD}}^2).$$

$$S_P(\alpha) \equiv S_{n_0}(\alpha), \quad \text{where} \quad S_n(\alpha) = \sum_{j=0}^n p_j \alpha^{j+1}.$$

$n_0 \equiv n_0(\alpha)$ is the order for which $p_{n_0} \alpha^{n_0+1}$ is minimal.




Figure : $\langle P \rangle_{MC}(\alpha) - S_n(\alpha)$ between MC data and sums truncated at orders α^{n+1} ($S_{-1} = 0$) vs. $a(\alpha)/r_0$. The lines $\propto a^i$ are drawn to guide the eye.




Figure : $\langle P \rangle_{\text{MC}}(\alpha) - S_P(\alpha)$. The linear fit is to $a^4 < 0.0013 r_0^4$ points only.




Figure : $\langle G^2 \rangle$ evaluated using the $N = 16$ and $N = 32$ MC data of Boyd et al. The error band is our prediction for $\langle G^2 \rangle$.

$$\langle G^2 \rangle = 3.18(29)r_0^{-4} = 24.2(8.0)\Lambda_{\overline{\text{MS}}}^4 \simeq 0.077 \text{ GeV}^4.$$




Figure : $aE_{MC} - a\delta m$ vs. a/r_0 . The expansion of $a\delta m$ was also converted into the \overline{MS} scheme at two (\overline{MS}_2) and three (\overline{MS}_3) loops. The curves are fits to $\bar{\Lambda}a + ca^2$.

Determination of N_m

$$\nu \sim m$$

Large β_0 analysis

$$m \left(\frac{\nu}{m} \right)^{2u} \simeq \nu \{ 1 + (2u - 1) \ln \frac{\nu}{m} + \dots \}.$$

Therefore, the underlying assumption is that we are in a regime where (besides $2u - 1 \ll 1$)

$$(2u - 1) \ln \frac{\nu}{m} \ll 1.$$

$$N_m = \frac{r_n}{(r_n^{\text{asym}} / N_m)}$$

Determination of N_m

$$\nu \sim m$$

Large β_0 analysis

$$m \left(\frac{\nu}{m} \right)^{2u} \simeq \nu \{ 1 + (2u - 1) \ln \frac{\nu}{m} + \dots \}.$$

Therefore, the underlying assumption is that we are in a regime where (besides $2u - 1 \ll 1$)

$$(2u - 1) \ln \frac{\nu}{m} \ll 1.$$

$$N_m = \frac{r_n}{(r_n^{\text{asym}} / N_m)}$$




Figure : N_m for $n_f = 3$, as a function of $x \equiv \mu/m_b$, obtained from r_n/r_n^{asym} with r_n^{asym} truncated at $\mathcal{O}(1/n^3)$. We name the different lines as NLO (dashed-dotted), NLO (dashed) and NNLO (solid) for $n = 0, 1, 2$, respectively.

The static potential

$$V(r; \nu_{us}) = \sum_{n=0}^{\infty} V_n \alpha_s^{n+1},$$

$2m_{\text{OS}} + V$ can be understood as an observable up to $\mathcal{O}(r^2 \Lambda_{\text{QCD}}^3, \Lambda_{\text{QCD}}^2/m)$ contributions $\rightarrow 2N_m + N_V = 0$ and

$$V_n^{\text{asym}} = N_V \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right)$$

$$N_V = \frac{V_n}{(V_n^{\text{asym}} / N_V)}$$

The static potential

$$V(r; \nu_{us}) = \sum_{n=0}^{\infty} V_n \alpha_s^{n+1},$$

$2m_{\text{OS}} + V$ can be understood as an observable up to $O(r^2 \Lambda_{\text{QCD}}^3, \Lambda_{\text{QCD}}^2/m)$
contributions $\rightarrow 2N_m + N_V = 0$ and

$$V_n^{\text{asym}} = N_V \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right)$$

$$N_V = \frac{V_n}{(V_n^{\text{asym}}/N_V)}$$

The static potential

$$V(r; \nu_{us}) = \sum_{n=0}^{\infty} V_n \alpha_s^{n+1},$$

$2m_{\text{OS}} + V$ can be understood as an observable up to $O(r^2 \Lambda_{\text{QCD}}^3, \Lambda_{\text{QCD}}^2/m)$
contributions $\rightarrow 2N_m + N_V = 0$ and

$$V_n^{\text{asym}} = N_V \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right)$$

$$N_V = \frac{V_n}{(V_n^{\text{asym}}/N_V)}$$




Figure : $-N_V/2 = N_m$ for $n_l = 3$, as a function of $x \equiv \nu r$, obtained from $-(N_V/2)v_n/v_n^{\text{asym}}$. v_n^{asym} is truncated at $\mathcal{O}(1/n^3)$.

$$N_m(n_l = 0) = 0.600(29)$$

$$N_m(n_l = 3) = 0.563(26)$$

~ 20 standard deviations from zero!

Conformal window: n_l dependence




Figure : $N_m(x = 1)$ obtained from $-(N_V/2)v_3/v_3^{\text{asym}}$ as a function of n_l .

First numerical evidence of the disappearance of the renormalon in the conformal window.




Figure : $c_n^{(3,0)}(N_S)/c_n^{(3,0)} - 1$ for $n \in \{0, 1, 2, 3, 4, 5, 7, 9, 11, 15\}$ (top to bottom). For each value of N_S we have plotted the data point with the maximum value of N_T . The curves represent the global fit. $-(1/N_S)f_{0,\text{DLPT}}^{(3,0)}/c_{0,\text{DLPT}}^{(3,0)}$ is shown for $n = 0$.




Figure : Zoom of previous Figure for $n = 9$.