Wojciech Flieger

U. Silesia in collaboration with Krzysztof Bielas & Janusz Gluza (U. Silesia), Marek Gluza (FU Berlin)

Singular values, contractions, dilations and neutrino mixing analysis

arXiv:1708.09196

Podlesice

3-8.09.2017

What is measured?



Wojciech Flieger |

Neutrino mixing in the Standard Model

$$\nu_{\alpha}^{(f)} = (U_{PMNS})_{\alpha i} \nu_{i}^{(m)}$$

Mixing matrix

$$U_{PMNS} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Experimental values of mixing parameters

$$\begin{array}{ll} \theta_{12} \in [31.38^{\circ}, 35.99^{\circ}], & \theta_{23} \in [38.4^{\circ}, 53.0^{\circ}], \\ \theta_{13} \in [7.99^{\circ}, 8.91^{\circ}], & \delta \in [0, 2\pi] \end{array}$$

Full experimental data - interval matrix

$$U_{PMNS} \xrightarrow{experiments} V_{osc}$$

CP Invariant Case

$$V_{osc} = \begin{pmatrix} 0.799 \div 0.845 & 0.514 \div 0.582 & 0.139 \div 0.155 \\ -0.538 \div -0.408 & 0.414 \div 0.624 & 0.615 \div 0.791 \\ 0.22 \div 0.402 & -0.73 \div -0.567 & 0.595 \div 0.776 \end{pmatrix}$$

http://www.nu-fit.org

$$V_{osc} \xrightarrow{?} BSM$$

Extended mixing - BSM models

Complete mixing

$$\left(\begin{array}{c}\nu^{(f)}\\\hat{\nu}^{(f)}\end{array}\right) = \left(\begin{array}{c}V_{osc}&V_{lh}\\V_{hl}&V_{hh}\end{array}\right) \left(\begin{array}{c}\nu^{(m)}\\\hat{\nu}^{(m)}\end{array}\right) \equiv U\left(\begin{array}{c}\nu^{(m)}\\\hat{\nu}^{(m)}\end{array}\right)$$

Observable part

$$\nu_{\alpha}^{(f)} = \underbrace{(V_{osc})_{\alpha i} \nu_{i}^{(m)}}_{\text{SM part}} + \underbrace{(V_{lh})_{\alpha j} \hat{\nu}_{j}^{(m)}}_{\text{BSM part}}$$

A standard approach to deviation from unitarity

$$\mathcal{M}_{PMNS}\mathcal{M}_{PMNS}^{\dagger} \equiv [(1+\eta)N][(1+\eta)N]^{\dagger} = 1 + \epsilon$$

 $N - \text{Unitary}$
 $\eta, \epsilon - \text{Hermitian}$

Our approach: mixing matrix and singular values

Singular values σ_i of a given matrix A are positive square roots of the eigenvalues λ_i of the matrix AA^{\dagger}

$$\sigma_i(A) = \sqrt{\lambda_i(AA^{\dagger})}$$

Properties:

- generalization of eigenvalues
- always positive
- stable under perturbations

Unitary matrices

 $UU^{\dagger} = I = diag(1, 1, ..., 1) \Longrightarrow$ all singular values equal to 1



Wojciech Flieger |





$\|A\| \leq 1$

Operator norm (spectral norm)

$$||A|| := \sup_{||x||=1} ||Ax|| = \sigma_{\max}(A)$$

Contractions as submatrices of the unitary matrix

$$\left\| \begin{pmatrix} \mathbf{V}_{osc} & V_{lh} \\ V_{hl} & V_{hh} \end{pmatrix} \right\| = 1 \Longrightarrow \| \mathbf{V}_{osc} \| \le 1$$

For U_{PMNS} holds

$$\sum_{\alpha} P_{\alpha\beta} = 1,$$

However, for a nonunitary U this relation is not fulfilled. $\Theta_2 = \Theta_1 + \varepsilon$

$$U = \begin{pmatrix} \cos \Theta_1 & \sin \Theta_1 \\ -\sin \Theta_2 & \cos \Theta_2 \end{pmatrix}$$

In this case we get, $\Delta_{ij} \propto (m_i^2 - m_j^2) rac{L}{E}$

$$P_{ee} + P_{e\mu} = 1 + 4\epsilon \sin^2 \Delta_{21} \sin \Theta_1 \cos \Theta_1 \cos 2\Theta_1 + \mathcal{O}(\epsilon^2)$$

$$P_{\mu e} + P_{\mu \mu} = 1 - 4\epsilon \sin^2 \Delta_{21} \sin \Theta_1 \cos \Theta_1 \cos 2\Theta_1 + \mathcal{O}(\epsilon^2)$$

Peculiar fact:

 $||U|| \ge 1$

Non-physical parametrization!

Statistics of Contractions in Vosc

Experimental mixing matrix

$$V_{osc} = \begin{pmatrix} 0.799 \div 0.845 & 0.514 \div 0.582 & 0.139 \div 0.155 \\ -0.538 \div -0.408 & 0.414 \div 0.624 & 0.615 \div 0.791 \\ 0.22 \div 0.402 & -0.73 \div -0.567 & 0.595 \div 0.776 \end{pmatrix}$$

Contractions: only 4 %

Non-physical: 96%

Contractions as a convex combination of unitary matrices

$$V = \sum_{i=1}^{m} \alpha_i U_i, \quad \alpha_i \ge 0 \text{ and } \sum_{i=1}^{m} \alpha_1 = 1$$
$$\|V\| = \|\sum_{i=1}^{m} \alpha_i U_i\| \le \sum_{i=1}^{m} \alpha_i \|U_i\| = 1$$

Physical Region





Wojciech Flieger

Unitary dilation

$U_{PMNS} ightarrow V_{osc} \xrightarrow{\text{contractions}} \Omega$

BSM?

Contractions

$$\boldsymbol{V} \in \Omega \xrightarrow{\text{dilation}} \begin{pmatrix} \boldsymbol{V} & V_{lh} \\ V_{hl} & V_{hh} \end{pmatrix} \equiv \boldsymbol{U} \rightarrow \boldsymbol{U}\boldsymbol{U}^{\dagger} = \boldsymbol{I}$$

CS decomposition

$$U \equiv \begin{pmatrix} \mathbf{V} & V_{lh} \\ V_{hl} & V_{hh} \end{pmatrix} = \begin{pmatrix} W_1 & 0 \\ 0 & W_2 \end{pmatrix} \begin{pmatrix} C & -S & 0 \\ S & C & 0 \\ 0 & 0 & I_{m-n} \end{pmatrix} \begin{pmatrix} Q_1^{\dagger} & 0 \\ 0 & Q_2^{\dagger} \end{pmatrix}$$

where $C \ge 0$ and $S \ge 0$ are diagonal matrices satisfying $C^2 + S^2 = I_n$ $W_1, Q_1 \in M_{n \times n}$ and $W_2, Q_2 \in M_{m \times m}$ are unitary matrices

12

As an illustration let us take two U_{PMNS} matrices

$$\begin{array}{ll} U_1: \ \theta_{12}=31.38^\circ, \theta_{23}=38.4^\circ, \theta_{13}=7.99^\circ, \\ U_2: \ \theta_{12}=35.99^\circ, \theta_{23}=52.8^\circ, \theta_{13}=8.90^\circ, \end{array}$$

and let us construct a contraction as

$$V = \frac{1}{2}U_1 + \frac{1}{2}U_2,$$

The set of singular values

$$\sigma_1(V) = 1, \ \sigma_2(V) = 0.991, \ \sigma_3(V) = 0.991$$

for which we get the following unitary dilation

	0.822411	0.548133	0.146854 0.70103	0.0169583	-0.0368511 0.0197681	
U = 1	0.311417	-0.643236	0.686702	0.0250273	0.130689	
	-0.0524981	0.122242	-0.0336064	0.599485	0.788536	-
	-0.0671638	0.00403263	0.119588	0.788536	-0.599485)
	\					

Quark Sector



Wolfenstein parametrization

$$s_{12} = \lambda, \quad s_{23} = A\lambda^2, \quad s_{13}e^{i\delta} = A\lambda^3(\rho + i\eta)$$
$$V_{CKM} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(\rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$

Distribution of contractions

All matrices within V_{CKM} are contractions with 2% accuracy

$$\begin{split} & \textbf{6\% of } \|V_{\rm CKM}\| = 1.002, \\ & \textbf{94\% of } \|V_{\rm CKM}\| = 1.001 \end{split}$$

$$0.961 \le \|V_{\rm osc}\| \le 1.178$$





- Interval matrix Vosc allows for independent analysis of mixing data
- Matrix theory and convex geometry offer suitable tools for that
- Singular values enrich studies beyond unitarity
- Contractions are natural to describe interplay between SM and BSM mixing theories in V_{osc}. They define physical region Ω by U_{PMNS} convex combination.
- ► There is a lot of space for BSM in *V*_{osc}. **Dilations** allow for appropriate construction of complete unitary matrices

Details in

A novel approach to neutrino mixing analysis based on singular values arXiv:1708.09196

Backup slides



Matrix norm



A matrix norm is a function $\|\cdot\|$ from the set of all complex (real matrices) into $\mathbb R$ that satisfies the following properties

$$\begin{split} \|\boldsymbol{A}\| &\geq \mathbf{0} \text{ and } \|\mathbf{A}\| = \mathbf{0} \Longleftrightarrow \mathbf{A} = \mathbf{0}, \\ \|\boldsymbol{\alpha}\boldsymbol{A}\| &= |\boldsymbol{\alpha}| \|\boldsymbol{A}\|, \boldsymbol{\alpha} \in \boldsymbol{C}, \\ \|\boldsymbol{A} + \boldsymbol{B}\| &\leq \|\boldsymbol{A}\| + \|\boldsymbol{B}\|, \\ \|\boldsymbol{A}\boldsymbol{B}\| &\leq \|\boldsymbol{A}\| \|\boldsymbol{B}\| \end{split}$$

Examples of matrix norms

- spectral norm: $||A|| = \max_{||x||_2=1} ||Ax||_2 = \sigma_1(A)$
- Frobenius norm: $||A||_F = \sqrt{Tr(A^{\dagger}A)} = \sqrt{\sum_{i,j=1}^n |a_{ij}|^2} = \sqrt{\sum_{i=1}^n \sigma_i^2}$
- ► maximum absolute column sum norm: ||A||₁ = max_{||x||1} ||Ax||_∞ = max_j ∑_j |a_{ij}|
- ► maximum absolute row sum norm: ||A||_∞ = max_{||x||∞}=1 ||Ax||_∞ = max_i ∑_i |a_{ij}|

Weyl's inequality for singular values

Let A and B be a $m \times n$ matrices and let $q = \min\{m, n\}$. Then

$\sigma_j(A+B) \leq \sigma_i(A) + \sigma_{j-i+1}(B)$ for $i \leq j$

Let us calculate UU^T and U^TU for U, $s(c)_i \equiv \sin(\cos)\Theta_i$, i = 1, 2

$$UU^{T} = \begin{pmatrix} 1 & s_{1}c_{2} - s_{2}c_{1} \\ s_{1}c_{2} - s_{2}c_{1} & 1 \end{pmatrix}$$
$$U^{T}U = \begin{pmatrix} c_{1}^{2} + s_{2}^{2} & c_{1}s_{1} - s_{2}c_{2} \\ c_{1}s_{1} - s_{2}c_{2} & s_{1}^{2} + c_{2}^{2} \end{pmatrix}$$

As for the real *A* we have $||A^TA|| = ||AA^T|| = ||A||^2$, we can focus only on one of this products. Let us then write UU^T in the following form

$$UU^{T} = \begin{pmatrix} 1 & s_{1}c_{2} - s_{2}c_{1} \\ s_{1}c_{2} - s_{2}c_{1} & 1 \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & s_{1}c_{2} - s_{2}c_{1} \\ s_{1}c_{2} - s_{2}c_{1} & 0 \end{pmatrix}$$

19

This can be simplified into

$$UU^{T} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & s_{3} \\ s_{3} & 0 \end{pmatrix} \equiv I + B$$

where $s_3 \equiv \sin \Theta_3 = \sin(\Theta_1 - \Theta_2)$.

Let us observe that eigenvalues of *B* are equal $\pm s_3$. Using fact that spectral norm is unitarily invariant and matrix *B* is symmetric, we get

$$||UU^{T}|| = ||I + B|| = ||W^{T}(I + B)W|| = ||I + W^{T}BW||$$

= ||I + D||

where W is an orthogonal matrix such that

$$W^T B W = D = diag(s_3, -s_3)$$



Since I + D equals

$$\left(\begin{array}{cc}1+s_3&0\\0&1-s_3\end{array}\right),$$

its operator norm, i.e., the largest singular value equals

$$1 + s_3$$
 if $s_3 \ge 0$,
 $1 - s_3$ if $s_3 < 0$

we can see that by adding *B* to identity matrix we can not decrease spectral norm

$$1 = \|I\| \le \|I + B\| = \|UU^T\|$$

Thus

$$||U|| \ge 1$$

Algorithm

The following steps lead to a contraction settled by U_{PMNS} and then to its unitary dilation of a minimal dimension

1) Select a finite number of unitary matrices U_i , i = 1, 2, ...m, within experimentally allowed range of parameters θ_{13} , θ_{23} and δ .

2) Construct a contraction U_{11} as a convex combination of selected matrices U_i

$$V = \sum_{i=1}^m \alpha_i U_i, \quad \alpha_1, \dots, \alpha_m \ge 0, \quad \sum_{i=1}^m \alpha_i = 1.$$

3) Find singular value decomposition of V, i.e.

$$V = W_1 \Sigma Q_1^{\dagger}$$

where W_1 , Q_1 are unitary, Σ is diagonal, and determine number η of singular values strictly less than 1.

4) Use CS decomposition

$$\begin{array}{ccc} U = \left(\begin{array}{ccc} V & V_{lh} \\ V_{hl} & V_{hh} \end{array} \right) = \\ \left(\begin{array}{ccc} W_1 & 0 \\ 0 & W_2 \end{array} \right) \left(\begin{array}{ccc} I_r & 0 & 0 \\ 0 & C & -S \\ \hline 0 & S & C \end{array} \right) \left(\begin{array}{ccc} Q_1^{\dagger} & 0 \\ 0 & Q_2^{\dagger} \end{array} \right) \end{array}$$

to find the unitary dilation $U \in \mathbb{M}_{(3+\eta) \times (3+\eta)}$ of contraction U_{11} .

Wojciech Flieger |